

Subscribers

Follow us

These findings, published in one of the world's leading scientific journals *Nature Medicine* in April, bring the scientists one step closer to developing a new way of tackling infections, by targeting microbial interactions rather than the specific microbes.

one another.

Assistant Professor Sanjay Haresh Chotirmall from the NTU Lee Kong Chian School of Medicine, who led the study, said, "Our current understanding of infections is that they occur when harmful microbes enter our bodies. This model of understanding, however, fails to account for resident microbes or explain why some patients with infection respond to antibiotics to which the microbe is resistant in laboratory testing. We are therefore proposing that microbes exist as networks, where interactions happen and that the resistant antibiotic in this case targets another microbe with which the culprit is interacting. We therefore can potentially improve clinical outcomes by breaking such crossfalk.

"The findings of our study are the first steps in providing a more holistic view of how infections occur. While our study looked at patients with bronchiectasis, we believe this concept applies to all forms of infection – whether skin, lung, or a gastrointestinal infection. This way of looking at infections potentially changes our understanding of infection and may offer fresh ways of treating them."

Associate Professor John Abisheganaden, co-author of the study and Head of Department of Respiratory & Critical Care Medicine at Tan Tock Seng Hospital, said: "By applying an integrated and holistic method, this study provides a new and fresh approach to our understanding of respiratory infection. Applying this precision-medicine approach can help the managing physician better understand and choose the most appropriate antibiotic or other therapy to confer clinical benefit – in short, to guide us to the right treatment at the right time, and for the best outcome."

Microbial interactions and infection

For their study, the scientists looked at patients with bronchiectasis, a disease of high Asian prevalence, where airways dilate irreversibly and, where infection promotes progression. Targeting bacteria with antibiotics reduces bacterial load and accompanying inflammation, which in turn alleviates symptoms and improves clinical outcomes.

To investigate interactions between microbes in the airways of patients with bronchiectasis, the team collected respiratory (sputum) samples from 383 patients from Singapore, Malaysia, Italy, and Scotland, including samples before, during and after bronchiectasis flare-ups.

After analysing the genetic material from bacteria, fungi and viruses in the samples, the scientists assessed for possible microbial interactions and found that patients with frequent flare-ups had more negative interactions, where microbes compete rather than cooperate, and that the number of such negative interactions increased even further during a flare-up. While changes to interactions between microbes during flare-ups were detected, there was surprisingly minimal change to the type and quantity of microbes present during a flare-up, and even after antibiotics were administered.

New treatment possibilities

The scientists believe that these findings suggest that microbial interactions potentially drive frequent flare-ups in patients.

Using these findings, the scientists have developed an online tool to help other researchers and physicians analyse microbial interactions in their own patient samples through the microbes' genetic sequences.

Asst Prof Chotirmall, who is also NTU Provost's Chair in Molecular Medicine, said: "We are proposing a fresh way to view infection, as networks rather than individual microbes. Targeting microbial interactions within an established network may promote more judicious antibiotic use and help curb rising antimicrobial resistance."

The team is currently exploring the use of probiotics to treat bronchiectasis by regulating microbiomes within the air passages.

The interdisciplinary team includes researchers from Singapore's NTU LKCMedicine, Tan Tock Seng Hospital, Changi General Hospital, and Singapore General Hospital; the University of Malaya in Malaysia, Italy's Fondazione IRCCCS Ca' Granda Ospedale Maggiore Policlinico and the University of Milan; the University of Sydney in Australia; the University of Dundee in Scotland; and the University of Exeter in the UK.

□ bacteria	bronchiectasis	fungi	gastrointestinal infection	lung infection	microbial interactions	NTU Singapore	
viruses							
□ Share > □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □							
*EXPRESS Pharma	EP News Bureau						
□ PREV POST						NEXT POST	

Unichem Labs gets US FDA approval for Amitriptyline HCL Tablets

Duke University researchers identify new coronavirus threat to humans