

# Science News

from research organizations

# Trial use of soybean waste to tackle obesity

Date:	March 22, 2022		
Source:	Nanyang Technological University		
Summary:	Scientists have found that fermented soybean waste, or okara, could improve fat metabolism and miti- gate effects of diet-induced obesity.		
Share:	f 🛩 🦻 in 🗷		
FULL STORY			

An international team of scientists from Nanyang Technological University, Singapore (NTU Singapore) and Waseda University in Japan have found that fermented soybean waste, or okara, could improve fat metabolism and mitigate effects of diet-induced obesity.

Through laboratory experiments reported in the peer-reviewed scientific journal *Metabolites* in February, the scientists from NTU Singapore and Waseda showed that mice on a high fat diet supplemented with fermented okara gained less body mass and had lower levels of fat and cholesterol after three weeks as compared to mice on the same diet but not fed any fermented okara.

With 14 million tonnes of okara generated every year, and nearly a third of the world's population overweight or obese, the scientists hope their findings can pave the way for fermented okara to be integrated into health foods one day, addressing the problems of food waste and obesity at the same time.

Dr Ken Lee, senior lecturer at the NTU School of Physical and Mathematical Sciences and co-lead author of the paper, said: "Using a process akin to what is used to produce miso and soy sauce, we fermented okara and fed it to mice on a high-fat diet. Three weeks later, we found that these mice gained less weight as compared to mice on a high-fat diet but without fermented okara. The mice that were fed fermented okara also had less fat and lower cholesterol levels.

"Our findings suggest that fermented okara could help to mitigate the harmful effects of a high-fat diet, and could potentially be integrated in the diets of obese patients who find it difficult to make immediate lifestyle changes."

Dr Shigenobu Shibata, Professor at the Waseda University Graduate School of Advanced Science and Engineering and co-lead author of the paper, said: "*Aspergillus oryzae* and *Aspergillus sojae*, whichare typical aspergillus fungi used to produce soy sauce and miso, were successfully combined to ferment okara. Compared to unfermented okara, fermented okara was found to be able to reduce obesity and improve abnormal lipid metabolism in mice. Tofu and soymilk are now widely accepted as health foods not only in Japan but in the world, and the effective use of okara, which is an industrial waste, is in line with the United Nations Sustainable Development Goals. I hope NTU and Waseda University, each with their unique strength, will continue to promote such joint research centred on graduate students and young researchers." The okara study supports the goals of the NTU 2025 five-year strategic plan, which prioritises the University's commitment to sustainability, as well as its focus on health and society as one of the research clusters with potential for significant intellectual and societal impact.

#### Enhancing okara's nutrition

To enhance the nutritional profile of okara, the research team from NTU and Waseda first fermented okara collected from a Singapore beancurd manufacturer using a process traditionally used in Asian countries to make fermented soybean food products such as miso, soy sauce, and tempeh.

Enhancing the nutritional profile of okara increases its potential as a desirable ingredient in the human diet, the scientists explained.

For this study, the team added a mix of *Aspergillus oryzae (A. oryzae)* and *Aspergillus sojae (A. sojae)* -- food-grade microbes -- to okara and left it to ferment in the dark at 25°C for four days.

Compared to unfermented okara, the scientists found that fermented okara had more protein and a higher total phenolic content -- an indication of higher antioxidant properties -- and less insoluble dietary fibre.

#### Fermented okara improves fat metabolism

The scientists then investigated the anti-obesity effects of fermented okara on mice put on a three-week high fat diet compared to mice put on three other types of diet: a normal diet, a high-fat diet, and a diet supplemented with unfermented okara.

At the end of three weeks, mice that were fed fermented okara every day gained the least body mass (3g) as compared to other three groups of mice that gained between 6g to 10g.

The mice that were fed fermented okara also had less visceral and subcutaneous fat (30g and 19g respectively) than mice on a high-fat diet without any fermented okara (67g and 53g respectively).

The scientists also found lower triglyceride and cholesterol levels in mice on a fermented okara-supplemented diet compared to those without. Triglycerides are a type of fat in the blood that has been linked to an increased risk of heart disease.

Further investigation revealed that the reduction in triglycerides was in part due to a significant dip in proteins involved in fatty acid synthesis.

These findings indicate the fermented okara's potential to suppress diet-induced obesity, said the scientists.

The study builds on the NTU-Waseda team's earlier research on other uses for okara. For instance, the scientists showed that fermented okara could help to bring down blood sugar levels. They have also worked with a local brewery to integrate okara into modern dishes.

Working with collaborators in Singapore, the NTU team has also successfully repurposed okara into an eco-friendly 3D-printing ink, highlighting the potential use of 3D food printing to improve food sustainability.

The NTU-Waseda team is now reaching out to industry partners to translate this work into healthy snacks and exploring how it can be integrated into other food products.

#### **Story Source:**

Materials provided by Nanyang Technological University. Note: Content may be edited for style and length.

#### **Related Multimedia:**

· YouTube video: NTUsg and Waseda University trial use of soybean waste to tackle obesity

#### Journal Reference:

 Natsumi Ichikawa, Li Shiuan Ng, Saneyuki Makino, Luo Lin Goh, Yun Jia Lim, Ferdinandus, Hiroyuki Sasaki, Shigenobu Shibata, Chi-Lik Ken Lee. Solid-State Fermented Okara with Aspergillus spp. Improves Lipid Metabolism and High-Fat Diet Induced Obesity. *Metabolites*, 2022; 12 (3): 198 DOI: 10.3390/metabo12030198

Cite This Page:	MLA	APA	Chicago

Nanyang Technological University. "Trial use of soybean waste to tackle obesity." ScienceDaily. ScienceDaily, 22 March 2022. <www.sciencedaily.com/releases/2022/03/220322150852.htm>.

# Featured Content

from New Scientist

Leaded petrol may have lowered the IQ of over half the US population

March 7, 2022 — Exposure to leaded petrol as a child has been linked to an average IQ drop of 2.6 points among US adults, increasing to 5.9 points among those born in the mid-to-late 1960s.

Hugging a pillow that mimics breathing could reduce anxiety

March 9, 2022 — Prototype pillow contains an inflatable chamber that connects to an external pump and motor, enabling it to expand and deflate like human lungs.

Electric field keeps kidney cells powered up while organs are on ice

March 9, 2022 — Organs soon run out of energy while they are between donor and recipient, but an electric field could keep them running and improve survival.

Visit New Scientist for more global science stories >>>

### RELATED STORIES

Shipment Tracking for 'Fat Parcels' in the Body

Oct. 14, 2019 — Without fat, nothing works in the body: These substances serve as energy suppliers and important building blocks -- including for the envelopes of living cells. Numerous diseases are related to ...

High-Fat Diets Affect Your Brain, Not Just Your Physical Appearance

Sep. 9, 2019 — Much research has pointed to how an unhealthy diet correlates to obesity, but has not explored how diet can bring about neurological changes in the brain. A recent study has discovered that high-fat ...

Caloric Restriction in Combination With Low-Fat Diet Helps Protect Aging Mouse Brains

Mar. 12, 2018 — New research finds that a low-fat diet in combination with limited caloric consumption prevents aginginduced inflammatory activation of immune cells in the mouse brain - and that exercise is ...

GM Soybean Oil Causes Less Obesity and Insulin Resistance but Is Harmful to Liver Function

Oct. 2, 2017 — Researchers have tested a genetically-modified soybean oil used in restaurants and found that while it induces less obesity and insulin resistance than conventional soybean oil, its effects on ...

## **Free Subscriptions**

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Email Newsletters

A RSS Feeds

### Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

- f Facebook
- ✓ Twitter
- in LinkedIn

## Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

- Leave Feedback
- Contact Us

About This Site | Staff | Reviews | Contribute | Advertise | Privacy Policy | Editorial Policy | Terms of Use

Copyright 2022 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners. Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.