

[Farming & Agriculture](#)[# Science & Research #](#)

Diversity in farm landscapes helps wildlife, global study finds

-- Greater variety within agricultural areas is a potential win-win for farmers and nature

Jun. 6, 2024

More diverse wildlife thrives on farms, benefitting both farmers and the environment, when there is variety in the agricultural landscape, a global study has found.



Farmland in China characterised by greater variability in the landscape, featuring a variety of crop types and semi-natural features, particularly along the borders of crop fields such as grassy strips. (Credit: NTU Singapore)

Specifically, the types of crops and non-crop features on farms, as well as how they are arranged, should be varied, according to research led by **Nanyang Technological University, Singapore (NTU Singapore)**.

The optimum approach would be to grow different crops like vegetables, cereals and fruit trees on the same farm in varying field sizes and layouts, as well as have semi-natural landscape features that are not crops, like shrubs, grassy borders and strips of wildflowers.

This variety increases the number of native animal and plant species such as pollinators and predators of pests, compared to agricultural lands that have just one type of crop grown on a single, uniform landscape.

Such an improvement in biodiversity arises because variability in crops and non-crop features, creates diverse "mini habitats" for different types of wildlife.

One of the main drivers of declines in biodiversity is increased farming to raise food production, so boosting the diversity of wildlife on farmlands is key in helping the environment.

Farmers stand to gain, too, as past research suggests that better farm biodiversity could improve crop yields; for example, by encouraging more pollinators of the crops and more predators to feed on crop pests, say the researchers of the latest study published in March 2024 in Ecology Letters.

Associate Professor Eleanor Slade, from NTU's Asian School of the Environment (ASE) and senior author of the study, said: "Our study demonstrates that maintaining diverse crop and non-crop habitats within agricultural landscapes is important for agricultural biodiversity on a global scale. This finding holds significant implications and provides valuable insights for policymakers, land managers and conservationists. By prioritising strategies that promote crop and landscape heterogeneity, stakeholders can help preserve biodiversity and also support sustainable agricultural practices."

The scientists also believe that increasing the diversity of crop types and non-crop features within farms could be a more appealing and practical approach for farmers to contribute to biodiversity conservation than existing methods.

Mr Tharaka S. Priyadarshana, a PhD student from ASE and the lead author of the study, said some strategies suggested that farmers reduce farming activity to restore semi-natural habitats that encourage wildlife, or consider organic farming that uses less chemicals.

"But when we look at the practicality of these strategies, they cannot be applied by many farmers, compared to having a good mix in crops and non-crop landscape features," he said.

Professor Emily Poppenborg Martin, from Germany's Justus Liebig University Giessen and the second author of the research, said: "This study is very important because of the crucial results it shows for a wide range of organisms, such as invertebrate predators and pollinators, which play key roles in agriculture. The evidence is clear: maintaining the biodiversity support system for the production of food, energy and feed goes hand-in-hand with learning how to diversify farmed landscapes globally."

Wide study scope



Farmland in China characterised by greater crop diversity, featuring a variety of crop types in smaller fields. (Credit: NTU Singapore)

The researchers analysed close to 6,400 farmlands reported in 122 previous studies conducted in 24 countries across Asia, Europe, North and South America. The wide scope of the study means its findings are more globally applicable than earlier studies, which drew conclusions largely based on North American and European countries.

The findings are relevant to a wider range of wildlife, too. Unlike past research that focussed only on specific types of animals, this study analysed many different types of wildlife including pollinators like bees, hoverflies and butterflies; predators like birds, carabid beetles and spiders; pests like aphids, plant bugs and weevils; herbaceous plants that do not have woody stems; and decomposers like dung beetles and springtails.

By analysing all the results from the earlier studies together, the scientists found that variety in crops and non-crop landscape elements on farms consistently improved the biodiversity of vertebrates like birds and invertebrates like insects, herbaceous plants, pollinators and predators.

Some types of wildlife had more gains in biodiversity than others. Statistically, vertebrates and plants benefitted more when both crop and non-crop features were diverse, while invertebrates showed similar benefits from either the diversification of crops or other non-crop features.

Pollinators gained more from having crops and non-crop elements with more varied layouts that better link them to one another, while predators favoured higher diversity within the types of crops and non-crop features.

These positive effects on farmlands were also consistently demonstrated in both tropical and temperate environments, in annual and perennial crops, and at different spatial scales, from within a few hundred metres of the farm to an area over several kilometres beyond the farm.

Why having diverse 'mini habitats' helps



A native Carpenter bee pollinating an eggplant in a farmland in Southern China, where woody semi-natural habitats are present. (Credit: NTU Singapore)

Mr Priyadarshana explained that growing diverse crop types alongside non-crop features on farms results in smaller areas for each crop type and creates mini habitats between different crops and non-crop fields. This encourages the presence of certain types of wildlife beneficial for crops.

For example, certain semi-natural features between crop borders, like grass strips with wildflowers, might attract certain insect pollinators that could help pollinate nearby crops. Certain crop types and semi-natural features could also attract specific predators that feed on the crop pests.

Mr Priyadarshana said the research findings could help balance farmers' goals to earn profits with sustainability and environmental aims.

"Agricultural systems are often managed to maximise their production in the economic sense. But by adapting farming practices and rethinking the way we integrate crop and non-crop habitats, they can also be managed to minimise environmental issues and maximise important services like pollination and pest control that boost crop production," he added.

The researchers now plan to look into the social aspects that need to be considered when trying to encourage farmers to introduce more diversity to the farms' crops and landscapes. For instance, more studies could be done to understand better what motivates farmers to plant a variety of crops compared to just one type of crop.

Paper titled "[Crop and landscape heterogeneity increase biodiversity in agricultural landscapes: A global review and meta-analysis](#)", published in Ecology Letters, 28 March 2024.

DOI: 10.1111/ele.14412

Source: [NTU Singapore](#)