



Search... Q

Login

Home > NEWS > Science News > Biology

# Singapore scientists pinpoint how non-alcoholic fatty liver disease increases risk of vascular diseases

BY BIOENGINEER — May 25, 2022 in Biology

0

Share on Facebook

Share on Twitter

Share on LinkedIn

Share on Reddit

Share on Telegram



These findings by scientists from **Nanyang Technological University, Singapore (NTU Singapore)** and the **National University of Singapore (NUS)** shed light on why the leading cause of mortality in NAFLD patients is cardiovascular complications, instead of liver damage.





					-25%	
		▲				
		▼				
			MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)	MEN AIRism Ultra Seamless Boxer...
			\$49.90	\$7.90	\$59.90	\$14.90

It is estimated that up to 40 per cent of the population in Southeast Asian countries, including Singapore, have NAFLD.

The research team, led by **Nanyang Assistant Professor Christine Cheung** from **NTU's Lee Kong Chian School of Medicine (LKCMedicine)**, found a higher level of blood vessel damage in NAFLD patients, which increases the risks of blood clots and cardiovascular diseases.

**Asst Prof Cheung** said: "The growing prevalence of fatty liver disease globally is a concern. In Singapore, one in three is likely to develop non-alcoholic fatty liver disease over the next 10 years. These patients are at increased risk of developing vascular diseases, such as coronary artery disease and cerebrovascular disease. The good news is that liver disease, at its early stages, is reversible."

AdChoices 

**EMERGENCY**

Built for  
Speed and  
Scale.

GET STARTED →



				-25%
	▲			
	▼	MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)
		\$49.90	\$7.90	\$59.90
				MEN AIRism Ultra Seamless Boxer...
				\$14.90

## Current Workforce Trends

Discover What Collaborative Communication Approaches Can Help Increase Innovation.

The team, comprising researchers from NTU, the National University Health System and the Agency for Science, Technology and Research's (A\*STAR), Singapore, published their findings in *EMBO Reports* in April.

### Fatty liver disease associated with increased chemokine levels

The team recreated blood vessel cells from samples donated by 99 NAFLD patients and 56 healthy controls to be used as experimental models for the study.



					-25%	
		▲				
		▼				
			MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)	MEN AIRism Ultra Seamless Boxer...
			\$49.90	\$7.90	\$59.90	\$14.90

## Download Our "2022 Global Workforce Trends" eBook

They found that the blood vessel cells of fatty liver disease patients contained higher levels of a class of proteins called **chemokines** – up to three times higher than in healthy individuals.

Chemokines are strong attractants of immune cells. Their main role in the body is drawing immune cells to sites of infections so that these cells can combat infections.

However, when abnormally high levels of immune cells are recruited into a blood vessel, it becomes harmful.

In particular, the researchers noticed that a type of immune cells known as **T cells** were strongly responsive to the chemokines in NAFLD patients. T cells are an important part of the body's immune system that fight against foreign particles.

The researchers discovered that the higher level of chemokines in fatty liver disease patients was attracting T cells into blood vessel walls. These T cells then caused inflammation of blood vessels, damaging them.



					-25%	
		▲				
		▼				
			MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)	MEN AIRism Ultra Seamless Boxer...
			\$49.90	\$7.90	\$59.90	\$14.90

Higher levels of damage and inflammation can lead to leakiness in the blood vessels, impairing blood vessels' integrity over time and increasing the risk of blood clot formation. It can lead to devastating events if the blood flow to vital organs is obstructed, such as ischaemic tissue injury.

The team detected three times as many circulating damaged endothelial cells — cells from the inner lining of blood vessels that have been shed into the bloodstream — in fatty liver disease patients, which is a sign of blood vessel injury.

The researchers believe that fatty liver disease patients have increased chemokine production due to the higher levels of oxidised lipids and free fatty acids in their blood.

The paper's first author, **NTU LKCMedicine research fellow Dr Ng Chun-Yi**, said: "Blood vessels are likely to be sensitive to the inflammatory mediators and abnormal lipid metabolism which underlie non-alcoholic fatty liver disease. We discovered that non-alcoholic fatty liver disease blood vessel cells are more 'activated', making them susceptible to vascular inflammation."

					-25%	
		▲				
		▼				
			MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)	MEN AIRism Ultra Seamless Boxer...
			\$49.90	\$7.90	\$59.90	\$14.90

Clinical collaborator and co-author **Associate Professor Dan Yock Young** from the **NUS Yong Loo Lin School of Medicine and the National University Health System** said: “Non-alcoholic fatty liver disease goes beyond the liver, impacting multi-organ system and leading to vascular complications such as coronary diseases and stroke. This study demonstrates for the first time how the levels of circulating damaged endothelial cells differ between non-alcoholic fatty liver disease patients and non-patients.”

“This opens new perspectives in our understanding of fatty liver diseases and therapeutic strategies for treating them. Instead of focusing narrowly on the liver pathology, a holistic approach in targeting the systemic inflammation of injured blood vessels has the potential to decrease the mortality rate of fatty liver disease patients,” said **Assoc Prof Dan**.

### Informing better clinical care for patients

Understanding how NAFLD affects heart health should help to inform clinical care for patients, the researchers say.

Typically, patients who present with symptoms of fatty liver diseases are not commonly referred to cardiologists. However, as the team’s research has shown that chronic



				-25%
	▲			
	▼	MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)
		\$49.90	\$7.90	\$59.90
				MEN AIRism Ultra Seamless Boxer...
				\$14.90

**Asst Prof Cheung** said: “As we better understand the link between non-alcoholic fatty liver disease and vascular inflammation, the science helps to inform clinical care. Patients who seek medical treatment for liver disease, for instance, could be considered for vascular health check-ups. This way, doctors not only follow up on the liver health but potentially prevent vascular complications in the future.”

“Liver disease at its early stage is reversible with lifestyle changes such as exercise and a healthy diet. Lifestyle modification is key at an early stage to slow down the progression of both fatty liver disease and cardiovascular disease before they become too advanced for functional recovery,” added Asst Prof Cheung.

Speaking independently on the study, **Professor Roger Foo, Zayed Bin Sultan Al Nahyan Professor in Medicine and Director of the Cardiovascular Disease Translational Research Programme at the NUS Yong Loo Lin School of Medicine and Senior Consultant at the National University Heart Centre, Singapore** said: “Non-alcoholic fatty liver disease is serious because if left untreated can cause significant and debilitating disease in the heart and walls of blood vessels. This work is important because, using samples donated from patients, the scientists have been able to map out new aspects of cellular crosstalk between cells of the vessel wall and the immune system, leading to a proposed fresh angle for treating the condition.”



				-25%
	▲			
	▼	MEN Slim Fit Chino Pants	MEN Woven Broadcloth Trunks	MEN Comfort Jacket (Pattern)
		\$49.90	\$7.90	\$59.90
				MEN AIRism Ultra Seamless Boxer...
				\$14.90

**Notes to Editor:**

Paper titled “*Endothelial-immune crosstalk contributes to vasculopathy in nonalcoholic fatty liver disease*”, published in *EMBO Reports*, 11 April 2022. DOI: 10.15252/embr.202154271

\*\*\*END\*\*\*

**Media contact:**

Mr Lester Hio  
 Manager, Media Relations  
 Corporate Communications Office  
 Nanyang Technological University, Singapore  
 Email: lester.hio@ntu.edu.sg

*About Nanyang Technological University, Singapore*

A research-intensive public university, Nanyang Technological University, Singapore (NTU Singapore) has 33,000 undergraduate and postgraduate students in the Engineering, Business, Science, Humanities, Arts, & Social Sciences, and Graduate colleges. It also has a medical school, the Lee Kong Chian School of Medicine, set up jointly with Imperial College London.

NTU is also home to world-class autonomous institutes – the National Institute of Education, S Rajaratnam School of International Studies, Earth Observatory of Singapore, and Singapore Centre for Environmental Life Sciences Engineering – and various leading research centres



					-25%	
		▲				
		▼				
			<b>MEN Slim Fit Chino Pants</b>	<b>MEN Woven Broadcloth Trunks</b>	<b>MEN Comfort Jacket (Pattern)</b>	<b>MEN AIRism Ultra Seamless Boxer...</b>
			\$49.90	\$7.90	\$59.90	\$14.90

Ranked amongst the world's top universities by QS, NTU has also been named the world's top young university since 2014. The University's main campus is frequently listed among the Top 15 most beautiful university campuses in the world and it has 57 Green Mark-certified (equivalent to LEED-certified) building projects, of which 95% are certified Green Mark Platinum. Apart from its main campus, NTU also has a campus in Singapore's healthcare district.

Under the NTU Smart Campus vision, the University harnesses the power of digital technology and tech-enabled solutions to support better learning and living experiences, the discovery of new knowledge, and the sustainability of resources.

For more information, visit [www.ntu.edu.sg](http://www.ntu.edu.sg)

## Journal

EMBO Reports

## DOI

10.15252/embr.202154271 [↗](#)

