Your source for the latest research news

Science News

from research organizations

Psychopathic individuals are more likely to have larger striatum region in the brain

Share:	f y 🤉 in 🖼
Summary:	Neuroscientists have established the existence of a biological difference between psychopaths and non-psychopaths.
Source:	Nanyang Technological University
Date:	May 10, 2022

FULL STORY

Neuroscientists from Nanyang Technological University, Singapore (NTU Singapore), University of Pennsylvania, and California State University, have established the existence of a biological difference between psychopaths and non-psychopaths.

Using magnetic resonance imaging (MRI) scans, they found that a region of the forebrain known as the striatum, was on average ten per cent larger in psychopathic individuals compared to a control group of individuals that had low or no psychopathic traits.

Psychopaths, or those with psychopathic traits, are generally defined as individuals that have an egocentric and antisocial personality. This is normally marked by a lack of remorse for their actions, a lack of empathy for others, and often criminal tendencies.

The striatum, which is a part of the forebrain, the subcortical region of the brain that contains the entire cerebrum, coordinates multiple aspects of cognition, including both motor and action planning, decision-making, motivation, reinforcement, and reward perception.

Previous studies have pointed to an overly active striatum in psychopaths but have not conclusively determined the impact of its size on behaviours. The new study reveals a significant biological difference between people who have psychopathic traits and those who do not.

While not all individuals with psychopathic traits end up breaking the law, and not all criminals meet the criteria for psychopathy, there is a marked correlation. There is clear evidence that psychopathy is linked to more violent behaviour.

The understanding of the role of biology in antisocial and criminal behaviour may help improve existing theories of behaviour, as well as inform policy and treatment options. To conduct their study, the neuroscientists scanned the brains of 120 participants in the United States and interviewed them using the Psychopathy Checklist -- Revised, a psychological assessment tool to determine the presence of psychopathic traits in individuals.

Assistant Professor Olivia Choy, from NTU's School of Social Sciences, a neurocriminologistwho co-authored the study, said: "Our study's results help advance our knowledge about what underlies antisocial behaviour such as psychopathy. We find that in addition to social environmental influences, it is important to consider that there can be differences in biology, in this case, the size of brain structures, between antisocial and non-antisocial individuals."

Professor Adrian Raine from the Departments of Criminology, Psychiatry, and Psychology at University of Pennsylvania, who co-authored the study, said: "Because biological traits, such as the size of one's striatum, can be inherited to child from parent, these findings give added support to neurodevelopmental perspectives of psychopathy -- that the brains of these offenders do not develop normally throughout childhood and adolescence."

Professor Robert Schug from the School of Criminology, Criminal Justice, and Emergency Management at California State University, Long Beach, who co-authored the study, said: "The use of the Psychopathy Checklist -- Revised in a community sample remains a novel scientific approach: Helping us understand psychopathic traits in individuals who are not in jails and prisons, but rather in those who walk among us each day."

Highlighting the significance of the work done by the joint research team, Associate Professor Andrea Glenn from the Department of Psychology of The University of Alabama, who is not involved in the research, said: "By replicating and extending prior work, this study increases our confidence that psychopathy is associated with structural differences in the striatum, a brain region that is important in a variety of processes important for cognitive and social functioning. Future studies will be needed to understand the factors that may contribute to these structural differences."

The results of the study were published recently in the peer-reviewed academic publication *Journal of Psychiatric Research*.

Bigger striatum, larger appetite for stimulation

Through analyses of the MRI scans and results from the interviews to screen for psychopathy, the researchers linked having a larger striatum to an increased need for stimulation, through thrills and excitement, and a higher likelihood of impulsive behaviours.

The striatum is part of the basal ganglia, which is made up of clusters of neurons deep in the centre of the brain. The basal ganglia receive signals from the cerebral cortex, which controls cognition, social behaviour, and discerning which sensory information warrants attention.

In the past two decades, however, the understanding of the striatum has expanded, yielding hints that the region is linked to difficulties in social behaviour.

Previous studies have not addressed whether striatal enlargement is observed in adult females with psychopathic traits.

The neuroscientists say that within their study of 120 individuals, they examined 12 females and observed, for the first time, that psychopathy was linked to an enlarged striatum in females, just as in males. In human development, the striatum typically becomes smaller as a child matures, suggesting that psychopathy could be related to differences in how the brain develops.

Asst Prof Choy added: "A better understanding of the striatum's development is still needed. Many factors are likely involved in why one individual is more likely to have psychopathic traits than another individual. Psychopathy can be linked to a structural abnormality in the brain that may be developmental in nature. At the same time, it is important to acknowledge that the environment can also have effects on the structure of the striatum."

Prof Raine added: "We have always known that psychopaths go to extreme lengths to seek out rewards, including criminal activities that involve property, sex, and drugs. We are now finding out a neurobiological underpinning of this impulsive and stimulating behaviour in the form of enlargement to the striatum, a key brain area involved in rewards.

The scientists hope to carry out further research to find out the causes of the enlargement of the striatum in individuals with psychopathic traits.

Story Source:

Materials provided by Nanyang Technological University. Note: Content may be edited for style and length.

Journal Reference:

1. Olivia Choy, Adrian Raine, Robert Schug. Larger striatal volume is associated with increased adult psychopathy. *Journal of Psychiatric Research*, 2022; 149: 185 DOI: 10.1016/j.jpsychires.2022.03.006

Cite This Page:

Nanyang Technological University. "Psychopathic individuals are more likely to have larger striatum region in the brain." ScienceDaily. ScienceDaily, 10 May 2022. <www.sciencedaily.com/releases/2022/05/220510102942.htm>.

Explore the latest scientific research on sleep and dreams in this free online course from New Scientist

--- Sign up now >>>

Featured Content

from New Scientist

Chicago

MLA

APA

Brain cells that are linked to Parkinson's disease finally identified

May 5, 2022 — High-tech genetic sequencing has found the brain cells connected to the movement disorder, which could lead to new treatments.

Vaccine may protect against the virus behind multiple sclerosis

May 4, 2022 — The jab could ward off Epstein-Barr virus, which causes glandular fever and is increasingly being linked to multiple sclerosis, lymphoma and stomach cancer.

How repealing Roe v Wade in the US will lead to more women's deaths

May 4, 2022 — A large body of evidence shows that restricting access to abortion doesn't reduce the number of abortions, only increases the risk of death for those who need them.

Visit New Scientist for more global science stories >>>

Explore More

from ScienceDaily

Not All Psychopaths Are Violent; A New Study May Explain Why Some Are 'Successful' Instead

May 12, 2020 — Psychopathy is widely recognized as a risk factor for violent behavior, but many psychopathic individuals refrain from antisocial or criminal acts. Understanding what leads these psychopaths to be ...

Psychopaths in the C-Suite?

Oct. 15, 2018 — The seemingly never-ending stream of corporate scandals over the past decades, from Enron to Theranos, suggests that something is rotten in corporate leaders. Many place the blame on psychopaths, who ...

Flourishing Under an Abusive Boss? You May Be a Psychopath, Study Shows

Jan. 23, 2018 — According to new research certain types of 'psychopaths' actually benefit and flourish under abusive ...

Neuroscientists Build Case for New Theory of Memory Formation

Oct. 23, 2017 — Neuroscientists propose that the existence of 'silent engrams' suggests current theories of memory formation need to ...

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Email Newsletters

A RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

- f Facebook
- ✓ Twitter
- in LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

- Leave Feedback
- Contact Us

Copyright 1995-2022 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners.

Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners. Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

- CCPA: Do Not Sell My Information - GDPR: Privacy Settings -