

# Computable Soft Separation Axioms

Authors

January 2022

## Abstract

Soft sets were introduced as a means to study objects that are not defined in an absolute way, and have found applications in numerous areas of mathematics, decision theory and in statistical applications. Soft topological spaces were first considered in [21], and soft separation axioms for soft topological spaces were studied in [6, 7, 4].

In this paper we introduce the effective versions of soft separation axioms. Specifically we focus our attention on computable u-soft and computable p-soft separation axioms and investigate various relations between them. We also compare the effective and classical versions of these soft separation axioms.

## 1 Preliminaries

### 1.1 Soft sets

Set theory classifies members of a set as whether those members belong to the set or not, but, in some situations we need to classify elements of a set based on some parameters. The need for such a classification motivated Molodtsov to introduce soft set theory in [14]. Soft set theory is considered a mathematical tool that deals with objects that are not defined in a definite way. Such objects can be found in complicated mathematical problems in economics and engineering applications when classical mathematical tools cannot be used due to the uncertainties associated to such problems. There are already existing mathematical tools for dealing with uncertainty in mathematical problems, such as the use of probability theory [10], fuzzy set theory [12] and interval mathematics [9]. However those three mathematical tools have their own shortcomings that the use of soft set theory overcomes as argued in [14].

Due to the unique properties of soft set theory that allow it to be more suitable in certain situations compared to the other mathematical tools mentioned above, it is often a major mathematical tool used in decision making problems as in [13] and [8]. Soft set theory, when combined with fuzzy set theory [12] can be used in decision making as in [28] and [18], and also used in forecasting problems as in [27]. There are also some applications of soft set theory in algebraic structures as in [1], [2] and [11]. When soft set theory is combined with rough set theory [16], we get new approximation spaces with interesting properties [20].

Topological spaces are introduced for soft sets [21], and some of the properties associated with soft topological spaces are explored in [15]. Several soft separation axioms were defined and studied in [6] and the further applications of those soft separation axioms are explored in [7] and [4]. Soft separation axioms are of importance in soft topological spaces as shown in the existing literature, much like how classical separation axioms have played a key role in the classification and the understanding of classical topological spaces. In this paper we will define and explore further soft separation axioms for soft topological spaces. We also define the computable versions of these soft separation axioms and investigate their properties in the effective setting. This paper is intended to investigate how computability interacts with soft topological spaces and soft separation axioms. Hence we will compare the various principles that arise by considering computable separation axioms in the soft setting.

The paper is organised as follows. In section 1.2 we recall some basic notions of soft sets and soft topological spaces as defined in the literature. In section 1.3 we briefly recall some notation and definitions that we will require from computable analysis, including computable topological spaces and computable separation axioms that were studied in the literature. In section 2 we define a new separation axiom for soft topological spaces, called *u-soft separation*, and give some of its basic properties. In section 3 we define and study computable u-soft separation axioms for computable soft spaces, and in section 4 we define and study various computable p-soft separation axioms. Finally in section 5 we compare the various principles introduced in sections 3 and 4.

## 1.2 Soft topological spaces

In this section, we recall some definitions and results of soft set theory and soft topological spaces. This section is meant to provide a self-contained introduction

to the basics and background of soft set theory. The initiated reader may skip ahead to section 1.3.

### 1.2.1 Basics of soft sets

**Definition 1.1.** [14] A pair  $(G, E)$  (usually denoted as  $G_E$ ) is called a soft set over a universe  $X$  if  $G$  is map from the parameter set  $E$  into  $2^X$ . We usually identify  $G_E = \{(e, G(e)) : e \in E \text{ and } G(e) \subseteq X\}$ .  $S(X_E)$  denotes the set of all soft sets over  $X$  with respect to the parameter set  $E$ . The relative complement of  $G_E$  is denoted by  $G_E^c$ , where  $G^c : E \rightarrow 2^X$  is defined by  $G^c(e) = X \setminus G(e)$ . Where the context is clear we do not refer to the universe  $X$ .

**Definition 1.2.** [13, 17] Soft union and soft intersection are taken parameter-wise. For two soft sets  $G_{E_1}, H_{E_2}$  over  $X$ , their soft union,  $G_{E_1} \cup H_{E_2}$ , is the soft set  $F_{E_1 \cup E_2}$  where  $F : E_1 \cup E_2 \rightarrow 2^X$  is defined as follows

$$F(e) = \begin{cases} G(e), & \text{if } e \in E_1 - E_2, \\ H(e), & \text{if } e \in E_2 - E_1, \\ G(e) \cup H(e), & \text{if } e \in E_1 \cap E_2. \end{cases}$$

The soft intersection  $G_{E_1} \cap H_{E_2}$  is the soft set  $I_{E_1 \cap E_2}$  where  $I(a) = G(a) \cap H(a)$  for every  $a \in E_1 \cap E_2$ .

Given  $x \in X$  and a soft set  $G_E$  there are four ways one can define membership or non-membership:

**Definition 1.3.** [14, 6] For a soft set  $G_E \in S(X_E)$  and  $x \in X$ , we say that

- $x \in G_E$  if  $x \in G(e)$  for each  $e \in E$ .
- $x \notin G_E$  if  $x \notin G(e)$  for some  $e \in E$ .
- $x \in G_E$  if  $x \in G(e)$  for some  $e \in E$ .
- $x \notin G_E$  if  $x \notin G(e)$  for each  $e \in E$ .

Hence  $\in$  and  $\notin$  are “strong” membership and non-membership respectively. Depending on the version of membership that one uses, the usual set theoretic operations might or might not be compatible:

**Proposition 1.4.** [6] For two soft sets  $G_E$  and  $H_E$  in  $S(X_E)$  and  $x \in X$ , we have the following,

1. If  $x \in G_E$ , then  $x \in G_E$ .
2.  $x \notin G_E$  if and only if  $x \in G_E^c$ .
3.  $x \in G_E \cup H_E$  if and only if  $x \in G_E$  or  $x \in H_E$ .
4. If  $x \in G_E \cap H_E$ , then  $x \in G_E$  and  $x \in H_E$ .
5. If  $x \in G_E$  or  $x \in H_E$ , then  $x \in G_E \cup H_E$ .
6.  $x \in G_E \cap H_E$  if and only if  $x \in G_E$  and  $x \in H_E$ .

**Definition 1.5.** [6, 13] A soft set  $G_E$  over  $X$  is said to be:

- A null soft set if  $G(e) = \emptyset$  for each  $e \in E$ . It is denoted by  $\tilde{\emptyset}$ .
- An absolute soft set if  $G(e) = X$  for each  $e \in E$ . It is denoted by  $\tilde{X}$ .
- A stable soft set if for some  $M \subseteq X$  we have  $G(e) = M$  for each  $e \in E$ .

There are two different ways one can define a point, either as a soft singleton or as a soft point:

**Definition 1.6.** [5, 21] The soft set  $x_E$  (called a soft singleton) is defined by  $x(e) = \{x\}$  for each  $e \in E$ . A soft point, denoted by  $p_e^x$ , is the soft set  $P_E$  where  $P(e) = \{x\}$  and  $P(k) = \emptyset$  for each  $k \in E \setminus \{e\}$ .

**Definition 1.7.** [17] A soft set  $G_{E_1}$  is a soft subset of a soft set  $H_{E_2}$ , denoted by  $G_{E_1} \subseteq H_{E_2}$ , if

- $E_1 \subseteq E_2$ , and
- $\forall e \in E_1, G(e) \subseteq H(e)$ .

Two soft sets are soft equal if each one of them is a soft subset of the other.

**Definition 1.8.** [19] The Cartesian product of two soft sets  $F_A$  and  $I_B$ , denoted by  $(F \times I)_{A \times B}$  over universes  $X$  and  $Y$ , respectively, is defined as  $(F \times I)(a, b) = F(a) \times I(b)$ , for each  $(a, b) \in A \times B$ .

### 1.2.2 Soft topological spaces

The study of soft topological spaces was initiated in [21]. We quickly recall some of the definitions and results of soft topological spaces.

**Definition 1.9.** [21, 15] A collection  $\tau$  of soft sets over a universe  $X$  w.r.t. a parameter set  $E$  is said to be a soft topology on  $X$  if the following conditions are satisfied,

1.  $\tilde{X}, \tilde{\emptyset} \in \tau$ .
2.  $\tau$  is closed under finite intersection.
3.  $\tau$  is closed under arbitrary union.

The triple  $(X, \tau, E)$  is called a *soft topological space*, or STS. Members of  $\tau$  are called soft open sets. A soft set is soft closed if its complement is soft open. The closure of  $H_E$ , denoted by  $\overline{H_E}$  is the intersection of all soft closed sets containing  $H_E$ .  $p_e^x$  is called a soft limit point of  $G_E$  if  $[F_E \setminus p_e^x] \cap G_E \neq \tilde{\emptyset}$ , for each soft open set  $F_E$  containing  $p_e^x$ .

Let  $Y$  be a nonempty soft subset of an STS  $(X, \tau, E)$ .  $\tau_Y = \{\tilde{Y} \cap G_E : G_E \in \tau\}$  is said to a soft relative topology on  $Y$  and the triple  $(Y, \tau_Y, E)$  is a soft subspace of  $(X, \tau, E)$ .

**Fact 1.10.** [21] Given an STS  $(X, \tau, E)$  and  $e \in E$ ,  $\tau_e = \{G(e) : G_E \in \tau\}$  forms a topology on  $X$  (classically).

**Theorem 1.11.** [19] Let  $(X, \tau, A)$  and  $(Y, \theta, B)$  be two STSs. Let  $\Omega = \{G_A \times F_B : G_A \in \tau \text{ and } F_B \in \theta\}$ . Then, the family of all arbitrary unions of elements of  $\Omega$  is a soft topology on  $X \times Y$ .

We now recall the partial soft separation axioms based on the partial membership ( $\in$ ) and strong non-membership ( $\notin$ ) relations:

**Definition 1.12.** [6] An STS  $(X, \tau, E)$  is said to be:

- p-soft  $T_0$  if for every two distinct  $x, y \in X$ , there exists a soft open set  $G_E$  such that  $x \in G_E$  and  $y \notin G_E$ , or  $y \in G_E$  and  $x \notin G_E$ .
- p-soft  $T_1$  if for every two distinct  $x, y \in X$ , there exist soft open sets  $G_E$  and  $F_E$  such that  $x \in G_E, y \notin G_E, y \in F_E$  and  $x \notin F_E$ .
- p-soft  $T_2$  if for every two distinct  $x, y \in X$ , there exist disjoint soft open sets  $G_E$  and  $F_E$  such that  $x \in G_E$  and  $y \in F_E$ .
- p-soft regular if for every soft closed set  $H_E$  and  $x \in X$  such that  $x \notin H_E$ , there exist disjoint soft open sets  $G_E$  and  $F_E$  such that  $H_E \subseteq G_E$  and  $x \in F_E$ .

Note that two soft sets are disjoint if their soft intersection is  $\tilde{\emptyset}$ .

The following well-know fact about  $T_1$  spaces holds in the p-soft setting:

**Theorem 1.13.** *An STS  $(X, \tau, E)$  is a p-soft  $T_1$  space if and only if  $x_E$  is soft closed, for all  $x \in X$ .*

### 1.3 Basics of computable analysis

#### 1.3.1 Type-2 theory of computability

Turing provided [22] in his pioneering work in 1936 an abstract model of a Turing machine. This is a central notion in the study of computability theory. In classical computability theory we deal with natural numbers and the domain and co-domain of computable functions are subsets of the natural numbers  $\mathbb{N}$ . However, in the study of effective analysis we are often concerned with potentially uncountable objects such as subsets of the real numbers, or sets of functions, etc. In order to apply the tools of classical computability we will need to “encode” these objects by means of *names*. Through systems of notations and representations in which the objects of study are represented as finite or infinite sequences of natural numbers we can make sense of the notion of a computation in which these names can be used as an input or the output of a computation.

Computable analysis has provided us with the formal framework in which we can conduct investigations of computability in the realm of analysis and topology. We introduce the notations that will be used throughout the paper. The reader is referred to [23, 25] for more details and background. Let  $\Sigma$  be a finite set of symbols that contains 0 and 1. The set of all finite words over  $\Sigma$  is denoted by  $\Sigma^*$ , and the set of all infinite sequences over  $\Sigma$  is denoted by  $\Sigma^\omega$  where  $q \in \Sigma^\omega$  means that  $q : \mathbb{N} \rightarrow \Sigma^\omega$  and we write  $q = q(0)q(1)\cdots$ , and  $|w|$  denotes the length of  $w \in \Sigma^*$ .  $q^{<i} \in \Sigma^*$  represents the initial segment of length  $i$  of  $q \in \Sigma^\omega$  and  $w \sqsubseteq q$  means that  $w$  is a prefix of  $q$ .

We use the wrapping function  $\iota : \Sigma^* \rightarrow \Sigma^*$ , where for  $a, b, c, d, e \in \Sigma$ ,  $\iota(abcde) = 110a0b0c0d0e011$  to encode the concatenation of finite strings in a way which can be effectively decoded. For instance, we cannot recover  $\sigma$  and  $\tau$  from  $\sigma\tau$  but we can do so from  $\iota(\sigma)\iota(\tau)$ . We fix the pairing function on the set of natural numbers as  $\langle i, j \rangle = \frac{(i+j)(i+j+1)}{2} + j$ . We also consider the standard tupling function on  $\Sigma^*$  and  $\Sigma^\omega$  where  $\langle v_1, \dots, v_n \rangle = \iota(v_1)\cdots\iota(v_n)$ ,  $\langle v, q \rangle = \iota(v)q$ ,  $\langle p, q \rangle = p(0)q(0)p(1)q(1)\cdots$ , and  $\langle q_0, q_1, \dots \rangle (\langle i, j \rangle) = q_i(j)$  for

$v_1, \dots, v_n, v \in \Sigma^*$  and  $p, q \in \Sigma^\omega$ . For  $r \in \Sigma^*$  let  $r^!$  be the longest subword  $s \in 11\Sigma^*11$  of  $r$  and then for  $u, r_1, r_2 \in \Sigma^*$ ,  $(u \ll r_1 \vee u \ll r_2) \Leftrightarrow u \ll r_1^! r_2^!$  where  $u \ll r$  iff  $\iota(u)$  is a subword of  $r$ .

For  $X_1, X_2 \in \{\Sigma^*, \Sigma^\omega\}$ , a (partial) function  $f : \subseteq X_1 \rightarrow X_2$  is computable if there is a type-2 machine  $M$  that computes  $f$  (see [23, 25] for more details if the reader is unfamiliar with the basics of effective type-2 theory). In TTE, we use representations or names to denote objects and type-2 machines can work with them via names. This is formalised through the notion of a represented space: a representation  $\delta$  of a set  $S$  is simply a surjective (partial) function  $\delta : \subseteq \Sigma^\omega \rightarrow S$ , while a notation  $\nu$  of a countable set  $S$  is a surjective (partial) function  $\nu : \subseteq \Sigma^* \rightarrow S$ . Examples include the canonical notations of the natural numbers and the rational numbers  $\nu_{\mathbb{N}} : \Sigma^* \rightarrow \mathbb{N}$ ,  $\nu_{\mathbb{Q}} : \Sigma^* \rightarrow \mathbb{Q}$ , respectively.

For representations or notations  $\gamma : \subseteq \Sigma^\omega \cup \Sigma^* \rightarrow M$  and  $\gamma' : \subseteq \Sigma^\omega \cup \Sigma^* \rightarrow M'$ , a partial function  $h : \subseteq \Sigma^\omega \cup \Sigma^* \rightarrow \Sigma^\omega \cup \Sigma^*$  realizes  $f : \subseteq M \rightarrow M'$  if  $f \circ \gamma(p) = \gamma' \circ h(p)$  for every  $p \in \text{dom}(\gamma)$ . The function  $f$  is called  $(\gamma, \gamma')$ -computable if it has a computable realization  $h$ . These definitions extend readily to multi-representations and multi-functions.

We say that  $\gamma$  is reducible to  $\gamma'$  (denoted by  $\gamma \leq \gamma'$ ) if  $M \subseteq M'$  and the identity function  $id : M \rightarrow M'$  is  $(\gamma, \gamma')$ -computable, i.e. there is a computable function that translates  $\gamma$ -names to  $\gamma'$ -names. Two representations  $\gamma$  and  $\gamma'$  are equivalent iff  $\gamma \leq \gamma'$  and  $\gamma' \leq \gamma$ .

Given a notation  $\alpha : \subseteq \Sigma^* \rightarrow M$  we can extend it naturally to a notation  $\alpha^{fs}$  for the set of finite subsets of  $M$ , and a representation  $\alpha^{cs}$  for the set of countable subsets of  $M$  in the natural way:

$$\alpha^{fs}(w) = W \Leftrightarrow (\forall u \ll w) u \in \text{dom}(\alpha), W = \{\alpha(u) : u \ll w\};$$

$$\alpha^{cs}(p) = W \Leftrightarrow (\forall u \ll p) u \in \text{dom}(\alpha), W = \{\alpha(u) : u \ll p\}.$$

If  $\mu : \subseteq \Sigma^\omega \rightarrow M'$  is a representation of  $M'$ , we can also define representations  $\mu^{fs}$  and  $\mu^{cs}$  for the set of finite and countable subsets of  $M'$  accordingly:  $\mu^{fs}(p) = W \Leftrightarrow (\exists n)(\exists q_1, \dots, q_n \in \text{dom}(\mu)), p = \langle 1^n, q_1, \dots, q_n \rangle, W = \{\mu(q_1), \dots, \mu(q_n)\}$ , and  $\mu^{cs}(\langle a_0 q_0, a_1 q_1, \dots \rangle) = W \Leftrightarrow (\forall i)(a_i = 0 \Rightarrow q_i \in \text{dom}(\mu))$  and  $W = \{\mu(q_i) : a_i = 0\}$ . Here  $w \in \Sigma^*$ ,  $p, q_0, q_1, \dots \in \Sigma^\omega$  and  $a_0, a_1, \dots$  are symbols of  $\Sigma$ .

### 1.3.2 Computable topological spaces

In this section, we define computable topological spaces as introduced in [24, 26] and mention some of the useful results in the literature that are relevant to us.

**Definition 1.14** (Weihrauch and Grubba [26]). An effective topological space is defined to be a 4-tuple  $\mathbf{X} = (X, \tau, \alpha, \mu)$  such that  $(X, \tau)$  is a topological  $T_0$  space and  $\mu : \subseteq \Sigma^* \rightarrow \alpha$  is a notation of a countable base  $\alpha$  of  $\tau$ .  $\mathbf{X}$  is a computable topological space if  $\text{dom}(\mu)$  is recursive and there is some c.e. set  $S$  such that for all  $u, v \in \text{dom}(\mu)$  we have

$$\mu(u) \cap \mu(v) = \bigcup \{ \mu(w) : (u, v, w) \in S \}.$$

In other words, the intersection of any two basic open sets is effectively open, uniformly in the notation for the basic open sets.

**Definition 1.15** (Weihrauch [24]). Let  $\mathbf{X} = (X, \tau, \alpha, \mu)$  be a computable topological space. We define the following representations.

1.  $\delta : \subseteq \Sigma^\omega \rightarrow X$  is a representation of the set  $X$ , where

$$\delta(p) = x \Leftrightarrow (\forall w \in \Sigma^*) (w \ll p \Leftrightarrow x \in \mu(w)).$$

2.  $\vartheta : \subseteq \Sigma^\omega \rightarrow \tau$  is a representation of the set of open sets where

$$\vartheta(p) = W \Leftrightarrow \forall w \in \Sigma^* (w \ll p \Rightarrow w \in \text{dom}(\mu)), \text{ and } W = \bigcup \{ \mu(w) : w \ll p \}. \blacksquare$$

3.  $\psi : \subseteq \Sigma^\omega \rightarrow \mathcal{A}$  is a representation of the set of closed sets where

$$\psi(p) = A \Leftrightarrow \forall w \in \Sigma^* (w \ll p \Leftrightarrow A \cap \mu(w) \neq \emptyset).$$

4.  $\varkappa : \subseteq \Sigma^\omega \rightrightarrows \mathbf{K}$  is a multi-representation of the set of compact subsets of  $X$  where

$$\varkappa(p) = K \Leftrightarrow [(\forall w \in \Sigma^*) (w \ll p \Leftrightarrow K \subseteq \cup \mu^{fs}(w))].$$

5.  $\bar{\delta} : \subseteq \Sigma^\omega \rightarrow X$  is a representation of the set  $X$ , where

$$\bar{\delta}(p) = x \Leftrightarrow \vartheta(p) = X \setminus \overline{\{x\}}.$$

6.  $\bar{\vartheta} : \subseteq \Sigma^\omega \rightarrow \tau$  is a representation of the set of open sets, where

$$\bar{\vartheta}(p) = X \setminus \psi(p).$$

7.  $\bar{\psi} : \subseteq \Sigma^\omega \rightarrow \mathcal{A}$  is a representation of the set of closed sets, where

$$\bar{\psi}(p) = X \setminus \vartheta(p).$$

We introduce some existing results that we will be using implicitly throughout the paper.

**Lemma 1.16** (Weihrauch [24]). *We have the following:*

1.  $\mu \leq \bigcup \mu^{fs} \leq \vartheta$ .
2.  $\delta(w \Sigma^\omega) = \bigcap \mu^{fs}(w)$  for all  $w \in \text{dom}(\mu^{fs})$ .
3. The space is  $SCT_2$  (see definition 1.19) iff  $\delta \leq \bar{\delta}$ .

The following theorem illustrates how we can compute unions and intersections of open and closed sets computably.

**Theorem 1.17** (Weihrauch [24]). *We have the following:*

1. Finite intersection on open sets is  $(\mu^{fs}, \vartheta)$ -computable and  $(\vartheta^{fs}, \vartheta)$ -computable.
2. Union on open sets is  $(\vartheta^{cs}, \vartheta)$ -computable.
3. Finite union on closed sets is  $(\bar{\psi}^{fs}, \bar{\psi})$ -computable, and intersection on closed sets is  $(\bar{\psi}^{cs}, \bar{\psi})$ -computable.
4. Finite union of compact sets is  $(\varkappa^{fs}, \varkappa)$ -computable.

**Lemma 1.18** (Weihrauch and Grubba [26]). *Given a point  $x$ , an open set  $W$ , a closed set  $A$  and a compact set  $K$ , we have the following:*

1. “ $x \in W$ ” is  $(\delta, \vartheta)$ -c.e.
2. “ $K \subseteq W$ ” is  $(\varkappa, \vartheta)$ -c.e.
3. “ $A \cap W \neq \emptyset$ ” is  $(\psi, \vartheta)$ -c.e.
4. “ $K \cap A = \emptyset$ ” is  $(\varkappa, \bar{\psi})$ -c.e.

### 1.3.3 Computable separation axioms

Weihrauch [24] introduced effective versions of separation axioms in computable topological spaces and discovered several interesting properties that hold for the computable separation axioms but not for their classical counterparts. For instance, he proved that the computable versions of  $T_2$  and  $T_1$  are equivalent [24] although they are clearly not classically equivalent.

In this section, we recall some of the computable separation axioms defined in [24] and the relationships between them. The main goal of this paper is to further this line of investigation for soft topological spaces. In the subsequent sections, we define different types of computable separation axioms for soft topological spaces and establish the relationships between them. We also show that certain implications are proper.

**Definition 1.19** (Weihrauch [24]). We define the following properties for a computable topological space  $(X, \tau, \alpha, \mu)$ :

- $CT_0$ : The multi-function  $t_0$  is  $(\delta, \delta, \mu)$ -computable, where  $t_0$  maps every pair of points  $(x, y) \in X^2$  such that  $x \neq y$  to some  $U \in \alpha$  such that  $x \in U$  and  $y \notin U$ , or  $x \notin U$  and  $y \in U$ .
- $CT_1$ : The multi-function  $t_1$  is  $(\delta, \delta, \mu)$ -computable, where  $t_1$  maps every pair of points  $(x, y) \in X^2$  such that  $x \neq y$  to some  $U \in \alpha$  such that  $x \in U$  and  $y \notin U$ .
- $CT_2$ : The multi-function  $t_2$  is  $(\delta, \delta, [\mu, \mu])$ -computable, where  $t_2$  maps every pair of points  $(x, y) \in X^2$  such that  $x \neq y$  to some  $(U, V) \in \alpha^2$  such that  $U \cap V = \emptyset$ ,  $x \in U$  and  $y \in V$ .
- $SCT_2$ : There is a c.e. set  $H \subseteq \Sigma^* \times \Sigma^*$  such that
  1.  $\forall x \neq y \exists (u, v) \in H (x \in \mu(u) \wedge y \in \mu(v))$ .
  2.  $\forall (u, v) \in H (\mu(u) \cap \mu(v) = \emptyset)$ .
- $CT_2^{pc}$ : The multi-function  $t^{pc}$  is  $(\delta, \varkappa, [\mu, \bigcup \mu^{fs}])$ -computable, where  $t^{pc}$  maps every  $x \in X$  and every compact set  $K$  such that  $x \notin K$  to some pair  $(U, W)$  of disjoint open sets such that  $x \in U$  and  $K \subseteq W$ .
- $CT_2^{cc}$ : The multi-function  $t^{cc}$  is  $(\varkappa, \varkappa, [\bigcup \mu^{fs}, \bigcup \mu^{fs}])$ -computable, where  $t^{cc}$  maps every pair  $(K, L)$  of non-empty disjoint compact sets to some pair  $(V, W)$  of disjoint open sets such that  $K \subseteq V$  and  $L \subseteq W$ .

- $SCT_2^{pc}$ : There is a c.e. set  $H \subseteq \Sigma^* \times \Sigma^*$  such that
  1.  $\forall x \in X \forall$  compact  $K$  such that  $x \notin K \exists (u, w) \in H (x \in \mu(u) \wedge K \subseteq \bigcup \mu^{fs}(w))$ .
  2.  $\forall (u, w) \in H (\mu(u) \cap \bigcup \mu^{fs}(w) = \emptyset)$ .
- $SCT_2^{cc}$ : There is a c.e. set  $H \subseteq \Sigma^* \times \Sigma^*$  such that
  1.  $\forall$  compact sets  $K, L$  such that  $K \cap L = \emptyset \exists (u, v) \in H (K \subseteq \bigcup \mu^{fs}(u)$  and  $L \subseteq \bigcup \mu^{fs}(v))$ .
  2.  $\forall (u, v) \in H (\bigcup \mu^{fs}(u) \cap \bigcup \mu^{fs}(v) = \emptyset)$ .

## 2 u-soft separation axioms

In section 1.2.2 we had mentioned soft separation axioms for STS based on strong membership and strong non-membership.

In this section, we define u-soft separation axioms. This type of separation axioms is based on soft points which is the natural way to define separation axioms analogously to the classical separation axioms. We investigate the relations between the u-soft separation axioms and p-soft separation axioms defined in [6]. we will note that some implications between the two different notions of soft separation axioms hold when the set of parameters is finite, however, when the parameter set is infinite those implications do not hold as what will be seen then from the counterexamples. We also answer a question proposed in [3] about whether u-soft  $T_2$  spaces imply p-soft  $T_2$  spaces where we find out that the answer is yes and we give a counterexample to show that the reverse implication is not true in general.

**Definition 2.1.** An STS  $(X, \tau, E)$  is called

- u-soft  $T_0$  iff  $\forall p_e^x, p_e^y \in \tilde{X}$ , there exists a soft open set  $G_E$  such that  $p_e^x \in G_E$  and  $p_e^y \notin G_E$ , or  $p_e^x \notin G_E$  and  $p_e^y \in G_E$ .
- u-soft  $T_1$  iff  $\forall p_e^x, p_e^y \in \tilde{X}$ , there exist two soft open sets  $G_E$  and  $F_E$  such that  $p_e^x \in G_E$  and  $p_e^y \notin G_E$ , and  $p_e^x \notin F_E$  and  $p_e^y \in F_E$ .
- u-soft  $T_2$  iff  $\forall p_e^x, p_e^y \in \tilde{X}$ , there exist two soft open sets  $G_E$  and  $F_E$  such that  $p_e^x \in G_E$  and  $p_e^y \in F_E$  and  $G_E \cap F_E = \tilde{\emptyset}$ .

Immediate implications between u-soft separation axioms are given in the next proposition.

**Proposition 2.2.** *Every u-soft  $T_i$  space is u-soft  $T_{i-1}$  space for  $i = 2, 1$ .*

*Proof.* Straightforward.  $\square$

Now, we give counterexamples to the above implications.

**Example 2.3.** *Let  $X = \{x\}, E = \{e_1, e_2\}$  and  $\tau = \{\tilde{X}, \tilde{\emptyset}, \{(e_1, \{x\}), (e_2, \tilde{\emptyset})\}\}$ .*

It can be easily seen that this space is u-soft  $T_0$  but not u-soft  $T_1$ .

**Example 2.4.** *Let  $E = \mathbb{N}, X$  be an infinite set,  $\tau = \{\tilde{X}, \tilde{\emptyset}, G_E : G_E^c \text{ is finite}\}$ .*

Clearly, this space is u-soft  $T_1$  but not u-soft  $T_2$ .

**Proposition 2.5.** *: An STS is a u-soft  $T_1$  space iff  $\forall p_e^x \in \tilde{X}, \overline{p_e^x} = p_e^x$ .*

*Proof.* Straightforward.  $\square$

The following propositions illustrate the relation between u-soft  $T_i$  and p-soft  $T_i$  spaces for  $i = 2, 1$ . Those implications are based on the finiteness of the parameter set and counterexamples are given to show that the implications are proper.

**Proposition 2.6.** *Every u-soft  $T_2$  space is p-soft  $T_2$  space if  $E$  is finite.*

*Proof.* Let  $x \neq y$  and  $E$  has  $m$  parameters.  $\forall p_{e_i}^x \forall p_{e_j}^y \in \tilde{X} \setminus p_{e_i}^x$ , there exist two disjoint soft open sets  $G_{E_{i_j}}$  and  $F_{E_{i_j}}$  such that  $p_{e_i}^x \in G_{E_{i_j}}$  and  $p_{e_j}^y \in F_{E_{i_j}}$ . Then,  $p_{e_i}^x \in \bigcap_{j=1}^m G_{E_{i_j}}$  and  $y \notin G_{E_{i_j}} \forall i \leq m$ , also,  $y \in \bigcup_{j=1}^m F_{E_{i_j}}$  and  $p_{e_i}^x \notin \bigcup_{j=1}^m F_{E_{i_j}}$ . Thus,

$$x \in \bigcup_{i=1}^m [\bigcap_{j=1}^m G_{E_{i_j}}] \text{ and } y \in \bigcap_{i=1}^m [\bigcup_{j=1}^m F_{E_{i_j}}],$$

and

$$[\bigcup_{i=1}^m [\bigcap_{j=1}^m G_{E_{i_j}}]] \cap [\bigcap_{i=1}^m [\bigcup_{j=1}^m F_{E_{i_j}}]] = \tilde{\emptyset}.$$

Therefore, the space is p-soft  $T_2$ .  $\square$

**Proposition 2.7.** *: Every u-soft  $T_1$  space is p-soft  $T_1$  space if  $E$  is finite.*

*Proof.* Let  $x \neq y$  and  $E$  has  $m$  parameters.  $\forall p_{e_i}^x \forall p_{e_j}^y \in \tilde{X} \setminus p_{e_i}^x$ , there exists an open set  $G_{E_{i_j}}$  such that  $p_{e_i}^x \in G_{E_{i_j}}$  and  $p_{e_j}^y \notin G_{E_{i_j}}$ . Then,  $p_{e_i}^x \in \bigcap_{j=1}^m G_{E_{i_j}}$ ,

$\forall i \leq m$ . Therefore,

$$x \in \bigcup_{i=1}^m [\bigcap_{j=1}^m G_{E_{i_j}}] \text{ and } y \notin \bigcup_{i=1}^m [\bigcap_{j=1}^m G_{E_{i_j}}].$$

Similarly, if we switch  $y$  and  $x$  we will get soft open sets  $G_{E_{i_j}}$  such that

$$y \in \bigcup_{i=1}^m [\bigcap_{j=1}^m F_{E_{i_j}}] \text{ and } x \notin \bigcup_{i=1}^m [\bigcap_{j=1}^m F_{E_{i_j}}].$$

Therefore, the space is p-soft  $T_1$ .  $\square$

The converse of the above propositions is not true in general as shown in the following example.

**Example 2.8.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  and  
 $\tau = \{\tilde{X}, \tilde{\emptyset}, \{(e_1, \{x\}), (e_2, \{x\})\}, \{(e_1, \{y\}), (e_2, \{y\})\}, \{(e_1, \emptyset), (e_2, \{x\})\},$   
 $\{(e_1, \{x\}), (e_2, \emptyset)\}, \{(e_1, \{y\}), (e_2, \emptyset)\}, \{(e_1, X), (e_2, \{x\})\}, \{(e_1, X), (e_2, \{y\})\},$   
 $\{(e_1, \{y\}), (e_2, X)\}, \{(e_1, X), (e_2, \emptyset)\}, \{(e_1, \{y\}), (e_2, \{x\})\}\}$ . This space is p-soft  $T_2$  but not u-soft  $T_1$ .

When the parameter set is infinite, the above inclusions do not hold in general as shown in the following examples.

**Example 2.9.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2, \dots\}$ . We define a STS  $\tau$  on  $X$  with respect to  $E$  as follows,  $\tau = \{\tilde{X}, \tilde{\emptyset}, G_k^{a_{i_1} a_{i_2} \dots a_{i_k}} : G_k^{a_{i_1} a_{i_2} \dots a_{i_k}} = \{(e_1, f(a_{i_1}), (e_2, f(a_{i_2}), \dots, (e_k, f(a_{i_k}), (e_{k+1}, X), \dots\}; i_1, \dots, i_k \in \{0, 1, 2, 3\}; f(a_0) = \emptyset, f(a_1) = \{x\}, f(a_2) = \{y\}, f(a_3) = X\}$ . Clearly, this space is u-soft  $T_1$  but it is not p-soft  $T_1$  or even p-soft  $T_0$ .

**Example 2.10.** Let  $X = \{a, b\}$ ,  $E = \{e_1, e_2, \dots\}$ . We partition  $\mathbb{N}$  into infinitely many infinite partitions  $\mathbb{N} = F_1 \cup F_2 \cup \dots$ . we define a STS on  $X$  with respect to  $E$  where its basic open sets are defined as follows, for each finite set  $G \subseteq \mathbb{N}$  we have  $\{p_{e_i}^a : i \in G\}$  and for each finite set  $G \subseteq \mathbb{N}$ ,  $n \in \mathbb{N}$  we have  $\{p_{e_n}^b\} \cup \{p_{e_i}^a : i \in F_n - G\}$ . Clearly, this space is u-soft  $T_2$  but it is not p-soft  $T_2$ .

The following two examples show that u-soft  $T_0$  and p-soft  $T_0$  spaces are incomparable.

**Example 2.11.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  and  
 $\tau = \{\tilde{X}, \tilde{\emptyset}, \{(e_1, \{y\}), (e_2, \{x\})\}, \{(e_1, \tilde{\emptyset}), (e_2, \{y\})\}, \{(e_1, \{y\}), (e_2, X)\},$   
 $\{(e_1, \tilde{\emptyset}), (e_2, \{x\})\}, \{(e_1, \tilde{\emptyset}), (e_2, X)\}\}$ .

This space is u-soft  $T_0$  but not p-soft  $T_0$ .

**Example 2.12.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$   
and  $\tau = \{\tilde{X}, \tilde{\emptyset}, \{(e_1, \{x\}), (e_2, \{x\})\}\}$ .

This space is p-soft  $T_0$  but not u-soft  $T_0$ .

### 3 Computable u-soft separation axioms

In this section, we define the new notions of computable soft topological spaces and computable u-soft separation axioms that are based on soft points. We investigate some properties and implications of those newly defined computable u-soft separation axioms. We also introduce some counterexamples to prove that some implications are not true in general.

**Definition 3.1.** A computable STS is a tuple  $(X, \tau, A, \beta, \nu)$  such that

1.  $(X, \tau, A)$  is a u-soft  $T_0$  space,
2.  $\nu : \Sigma^* \rightarrow \beta$  is a notation of a base of  $\tau$  with respect to soft points with recursive domain,
3. There is a computable function  $h : \Sigma^* \times \Sigma^* \rightarrow \Sigma^\omega$  such that for all  $u, v \in \text{dom}(\nu)$ ,

$$\nu(u) \bigcap \nu(v) = \cup \{\nu(w) : w \in \text{dom}(\nu) \text{ and } \iota(w) \ll h(u, v)\}.$$

**Note.** In computable soft topological spaces when we encode soft points we need to consider the parameter of the soft point so that it is encoded as well in the name. That is,  $\delta^u(p) = p_e^x$  where  $p$  is a list of all basic soft open sets containing  $p_e^x$  and the first bit of  $p$  encodes the parameter of the soft point, which is  $e$  in this case. When the parameter set  $E$  is infinite, we require it to be computable and countable, and to be given of the form  $E = \{e_1, e_2, \dots\}$ .

The following are the computable u-soft separation axioms where they are based on separating soft points by basic soft open sets.

**Definition 3.2.** A computable STS  $(X, \tau, A, \beta, \nu)$  is computable u-soft  $T_0$  ( $CuT_0$ , for short ) if  $(X, \tau, A)$  is a u-soft  $T_0$  and the multifunction  $ut_0$  is  $(\delta^u, \delta^u, \nu)$ -computable where  $ut_0$  maps every  $(p_e^x, p_\alpha^y) \in \tilde{X} \times \tilde{X}$  such that  $p_e^x \neq p_\alpha^y$  to some  $U_A \in \beta$  such that

$$(p_e^x \in U_A \text{ and } p_\alpha^y \notin U_A) \text{ or } (p_e^x \notin U_A \text{ and } p_\alpha^y \in U_A).$$

**Definition 3.3.** A computable STS  $(X, \tau, A, \beta, \nu)$  is computable u-soft  $T_1$  ( $CuT_1$ , for short ) if  $(X, \tau, A)$  is a u-soft  $T_0$  and the multifunction  $ut_1$  is  $(\delta^u, \delta^u, \nu)$ -computable where  $ut_1$  maps every  $(p_e^x, p_\alpha^y) \in \tilde{X} \times \tilde{X}$  such that  $p_e^x \neq p_\alpha^y$  to some  $U_A \in \beta$  such that

$$(p_e^x \in U_A \text{ and } p_\alpha^y \notin U_A).$$

**Definition 3.4.** A computable STS  $(X, \tau, A, \beta, \nu)$  is computable u-soft  $T_2$  ( $CuT_2$ , for short ) if  $(X, \tau, A)$  is a u-soft  $T_0$  and the multifunction  $ut_2$  is  $(\delta^u, \delta^u, \nu)$ -computable where  $ut_2$  maps every  $(p_e^x, p_\alpha^y) \in \tilde{X} \times \tilde{X}$  such that  $p_e^x \neq p_\alpha^y$  to some  $U_A, V_A \in \beta$  such that

$$(p_e^x \in U_A \text{ and } p_\alpha^y \in V_A \text{ and } U_A \cap V_A = \tilde{\emptyset}).$$

The next lemma gives the obvious implications between the computable u-soft separation axioms that are defined so far. The proof is Straightforward by definition.

**Lemma 3.5.**  $CuT_i \Rightarrow u\text{-soft}T_i$  for  $i \in \{0, 1, 2\}$ .

*Proof.* Straightforward. □

**Lemma 3.6.**  $CuT_i \Rightarrow CuT_{i-1}$  for  $i \in \{1, 2\}$ .

*Proof.* Straightforward. □

We give a counterexample that is  $CuT_0$  but not  $CuT_1$ .

**Example 3.7.** Let  $X = \{x\}$  be the universe set,  $E = \{e_1, e_2\}$  be a set of parameters and  $\tau$  is a STS generated by the following base,

$$\nu(01) = \{(e_1, \{x\}), (e_2, \emptyset)\}, \nu(001) = \tilde{X}, \text{ where } \beta = \text{range}(\nu).$$

We define now some more computable u-soft separation axioms to help us establish the relation between  $CuT_1$  and  $CuT_2$ . At the end of this section we will see that some of the following notions are equivalent.

**Definition 3.8.** A computable STS is:

$WCuT_0$ : If there is a c.e. set  $H \subseteq dom(\nu) \times dom(\nu)$  such that

1.  $(\forall p_e^x \neq p_\alpha^y)(\exists(u, v) \in H)(p_e^x \in \nu(u) \text{ and } p_\alpha^y \in \nu(v))$ ,
2.  $(\forall(u, v) \in H)$ :

$$\begin{aligned} &(\nu(u) \cap \nu(v) = \tilde{\emptyset}), \\ &\vee((\exists p_e^x)\nu(u) = \{p_e^x\} \subseteq \nu(v)), \\ &\vee((\exists p_\alpha^y)\nu(v) = \{p_\alpha^y\} \subseteq \nu(u)). \end{aligned}$$

$SCuT_0$ : If The multifunction  $ut_0^s$  is  $(\delta^u, \delta^u, [\nu_N, \nu])$ -computable where  $ut_0^s$  maps every  $(p_e^x, p_\alpha^y) \in \tilde{X} \times \tilde{X}$  such that  $(p_e^x \neq p_\alpha^y)$  to some  $(k, U_E) \in \mathbb{N} \times \beta$  such that

$$(k = 1, p_e^x \in U_E \text{ and } p_\alpha^y \notin U_E) \vee (k = 2, p_e^x \notin U_E \text{ and } p_\alpha^y \in U_E).$$

$CuT_0'$ : If there is a c.e. set  $H \subseteq dom(\nu_N) \times dom(\nu) \times dom(\nu)$  such that

1.  $(\forall p_e^x \neq p_\alpha^y)(\exists(w, u, v) \in H)(p_e^x \in \nu(u) \text{ and } p_\alpha^y \in \nu(v))$ ,
2.  $(\forall(w, u, v) \in H)$ :

$$\begin{aligned} &(\nu(u) \cap \nu(v) = \tilde{\emptyset}), \\ &\vee(\nu_N(w) = 1(\exists p_e^x)\nu(u) = \{p_e^x\} \subseteq \nu(v)), \\ &\vee(\nu_N(w) = 2(\exists p_\alpha^y)\nu(v) = \{p_\alpha^y\} \subseteq \nu(u)). \end{aligned}$$

$CuT_1'$ : If there is a c.e. set  $H \subseteq dom(\nu) \times dom(\nu)$  such that

1.  $(\forall p_e^x \neq p_\alpha^y)(\exists(u, v) \in H)(p_e^x \in \nu(u) \text{ and } p_\alpha^y \in \nu(v))$ ,
2.  $(\forall(u, v) \in H)$ :

$$\begin{aligned} &(\nu(u) \cap \nu(v) = \tilde{\emptyset}), \\ &\vee((\exists p_e^x)\nu(u) = \{p_e^x\} \subseteq \nu(v)). \end{aligned}$$

$CuT_2'$ : If there is a c.e. set  $H \subseteq dom(\nu) \times dom(\nu)$  such that

1.  $(\forall p_e^x \neq p_\alpha^y)(\exists(u, v) \in H)(p_e^x \in \nu(u) \text{ and } p_\alpha^y \in \nu(v))$ ,

2.  $(\forall(u, v) \in H)$ :

$$(\nu(u) \cap \nu(v) = \tilde{\emptyset}),$$

$$\vee((\exists p_e^x)\nu(u) = \{p_e^x\} = \nu(v)).$$

$SCuT_2$ : If there is a c.e. set  $H \subseteq \text{dom}(\nu) \times \text{dom}(\nu)$  such that

1.  $(\forall p_e^x \neq p_\alpha^y)(\exists(u, v) \in H)(p_e^x \in \nu(u) \text{ and } p_\alpha^y \in \nu(v))$ ,

2.  $(\forall(u, v) \in H)$ :

$$(\nu(u) \cap \nu(v) = \tilde{\emptyset}).$$

Now we investigate the relations between those separation axioms.

**Proposition 3.9.**  $CuT_0 \Leftrightarrow SCuT_0 \Leftrightarrow CuT_0'$ .

*Proof.*  $SCuT_0 \Rightarrow CuT_0$ : Straightforward.

$CuT_0 \Rightarrow SCuT_0$ : There is a machine  $M$  on input  $(p, q) \in \text{dom}(\delta^u) \times \text{dom}(\delta^u)$ , it first runs  $ut_0$  on  $(p, q)$  that outputs  $u$ . Then  $M$  outputs  $(1, u)$  if  $u \ll p$ , and outputs  $(2, u)$  if  $u \ll q$ .

$CuT_0' \Rightarrow SCuT_0$ : There is a machine  $M$  on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$ , it first searches for  $(w, u, v) \in H$  such that  $u \ll p$  and  $v \ll q$  and then it outputs  $(w, u)$  if  $\nu_N(w) = 1$  and  $(w, v)$ , otherwise.

$SCuT_0 \Rightarrow CuT_0'$ : Let  $M$  be a machine that realizes  $ut_0^s$ . There is another machine  $M'$  that on input  $(w, u, v) \in (\Sigma^*)^3$  halts iff we can find words  $u' \in \text{dom}(\nu)$ ,  $f, h \in \text{dom}(\nu^{fs})$  and  $t \leq \min(|f|, |h|)$  such that  $M$  on  $(f1^\omega, h1^\omega)$  halts in  $t$  steps outputting  $(w, u')$  and

$$u \ll g(f\nu(u')) \text{ and } v \ll g(h) \text{ if } \nu_N(w) = 1,$$

$$u \ll g(h) \text{ and } v \ll g(f\nu(u')) \text{ if } \nu_N(w) = 2.$$

Now, let  $H = \text{dom}(f_{M'})$ .

We need now to show the two conditions of  $H$ . For the first condition: Let  $\delta^u(p) = p_e^x \neq p_\alpha^y = \delta^u$ . Then  $M$  on  $(p, q)$  halts and outputs  $(w, u')$  in  $t$  steps where  $\nu_N(w) = 1, p_e^x \in \nu(u')$  and  $p_\alpha^y \notin \nu(u')$  (when  $\nu_N(w) = 2$ , same argument follows). Then,  $M$  also halts on  $(f1^\omega, h1^\omega)$  outputting  $(w, u')$  where  $f = p^{<t}$  and  $h = q^{<t}$ . Thus,  $p_e^x \in \cap \nu^{fs}(f\nu(u'))$  and  $p_\alpha^y \in \cap \nu^{fs}(h)$  and hence there are  $u, v$  such that  $u \ll \nu^{fs}(f\nu(u')), u \ll p$  and  $v \ll \nu^{fs}(h), v \ll q$ . Therefore, there exists some  $(w, u, v) \in H$  such that  $p_e^x \in \nu(u)$  and  $p_\alpha^y \in \nu(v)$ .

For the second condition of  $H$ : Let  $(w, u, v) \in H$ ,  $\nu_N(w) = 1$ ,  $p_e^x \in \nu(u)$ ,  $p_\alpha^y \in \nu(u) \cap \nu(v)$  and  $p_e^x \neq p_\alpha^y$ . Then, there are  $f, h, u'$  and  $t$  such that  $t \leq \min(|f|, |h|)$  and  $M$  halts on  $(f1^\omega, h1^\omega)$  in  $t$  steps outputting  $(w, u')$  and  $u \ll g(fu(u'))$  and  $v \ll g(h)$ . Therefore,  $p_e^x \in \nu(u) \subseteq \delta^u[f\Sigma^\omega] \cap \nu(u')$  and  $p_\alpha^y \in \nu(v) \subseteq \delta^u[h\Sigma^\omega]$ . We know that  $p_e^x \in \nu(u')$  and  $p_\alpha^y \notin \nu(u')$ . But,  $p_\alpha^y \in \nu(u) \subseteq \nu(u')$ , which a contradiction. Therefore, it must be the case that  $p_e^x = p_\alpha^y$ , hence,

$$((w, u, v) \in H, \nu_N(w) = 1 \text{ and } \nu(u) \cap \nu(v) \neq \tilde{\emptyset}) \Rightarrow (\exists p_e^x) \nu(u) = \{p_e^x\} \subseteq \nu(v).$$

Same argument follows when  $\nu_N(w) = 2$ . □

**Proposition 3.10.**  $SCuT_2 \Rightarrow CuT_2 \Rightarrow CuT_0 \Rightarrow WCuT_0$ .

*Proof.* Straightforward. □

**Proposition 3.11.**  $CuT_2 \Leftrightarrow CuT_2' \Leftrightarrow CuT_1 \Leftrightarrow CuT_1'$ .

*Proof.*  $CuT_1 \Leftrightarrow CuT_1'$ : Straightforward as it is a special case of  $SCuT_0 \Leftrightarrow CuT_0'$ .

$CuT_2' \Rightarrow CuT_1'$ : Straightforward.

$CuT_1' \Rightarrow CuT_2'$ : Let  $H$  be the c.e. set from  $CuT_1'$ . Now, let

$$H' = \{(r, s) : r \ll g(u, v'), s \ll g(u', v) \text{ for some } (u, v), (u', v') \in H\}.$$

We prove now the two conditions of  $H'$  as the c.e. set of  $CuT_2'$ .

Suppose  $p_e^x \neq p_\alpha^y$ . By the first condition of  $H$  there are  $(u, v), (u', v') \in H$  such that  $p_e^x \in \nu(u), p_\alpha^y \in \nu(v), p_\alpha^y \in \nu(u')$ , and  $p_e^x \in \nu(v')$ . Then,  $p_e^x \in \nu(u) \cap \nu(v')$  and  $p_\alpha^y \in \nu(u') \cap \nu(v)$ , and hence there is  $(r, s) \in H'$  such that  $p_e^x \in \nu(r)$  and  $p_\alpha^y \in \nu(s)$ . Thus the first condition of  $H'$  holds.

Now, we prove the second condition of  $H'$ . Suppose  $(r, s) \in H'$  and  $\nu(r) \cap \nu(s) \neq \tilde{\emptyset}$ . Thus, by definition of  $H'$  there are  $(u, v), (u', v') \in H$  such that  $\nu(r) \subseteq \nu(u) \cap \nu(v')$  and  $\nu(s) \subseteq \nu(u') \cap \nu(v)$ . Hence,  $\nu(u) \cap \nu(v) \neq \tilde{\emptyset}$  and  $\nu(u') \cap \nu(v') \neq \tilde{\emptyset}$ . Now, by the second condition of  $H$ ,  $\nu(u) = \{p_e^x\} \subseteq \nu(v)$  and  $\nu(u') = \{p_\alpha^y\} \subseteq \nu(v')$ . Therefore,  $\nu(r) = \{p_e^x\} = \nu(s)$  which shows that the second condition of  $H'$  holds.

$CuT_2' \Rightarrow CuT_2$ : There is a machine  $M$  that on input  $(p, q)$  searches for  $(u, v) \in H$  such that  $u \ll p$  and  $v \ll q$  and prints  $(u, v)$  if the search is successful and diverges, otherwise.

$CuT_2 \Rightarrow CuT_2'$ : By transitivity, which completes the proof. □

Now, we give counterexample to the above implications.

**Proposition 3.12.** *There is a STS that is  $WCuT_0$  but not  $CuT_0$ .*

*Proof.* Follows immediately from the next example.  $\square$

**Example 3.13.** *Let  $X = \{x_i, y_i : i \in \mathbb{N}\}$ ,  $E = \{e\}$  be a set of parameters, and  $\tau$  be the soft discrete topology defined on  $X$  w.r.t  $E$ . We will define  $A, B, C$ , and  $D$  as a partition of  $\mathbb{N}$  where  $A$  is a non c.e. set. We define a notation  $\nu$  of a basis of  $\tau$  as follows:*

	$0^i1$	$0^i2$	$0^i3$	$0^i12$	$0^i13$	$0^i23$
$i \in A \cup D$	$\{p_e^{x_i}\}$	$\{p_e^{y_i}\}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$
$i \in B$	$\{p_e^{x_i}\}$	$\{p_e^{x_i}, p_e^{y_i}\}$	$\{p_e^{y_i}\}$	$\{p_e^{x_i}\}$	$\tilde{\emptyset}$	$\{p_e^{y_i}\}$
$i \in C$	$\{p_e^{x_i}, p_e^{y_i}\}$	$\{p_e^{y_i}\}$	$\{p_e^{x_i}\}$	$\{p_e^{y_i}\}$	$\{p_e^{x_i}\}$	$\tilde{\emptyset}$

We define now the intersection of soft basic open sets computably,  $\nu(0^i m) \cap \nu(0^j n) = \nu(0^i m n)$  for  $m \neq n$ . Thus,  $(X, \tau, E, \beta, \nu)$  is a computable STS. Let  $H = \{(0^i m, 0^j n) : i, j \in \mathbb{N}; m, n \in \{1, 2\}; (i \neq j \text{ or } m \neq n)\}$ . Then  $H$  satisfies the two conditions of  $WuCT_0$ . We show now that this space is not  $SCuT_0$ . Let  $r, s \in \Sigma^{\mathbb{S}}$  such that  $\nu_N(r) = 1$  and  $\nu_N(s) = 2$ . W.L.O.G. assume that  $\nu_N$  is injective. For  $i \in \mathbb{N}$  let

$$S_i = \{\langle r, 0^i1 \rangle, \langle s, 0^i3 \rangle, \langle r, 0^i12 \rangle, \langle s, 0^i23 \rangle\},$$

$$T_i = \{\langle s, 0^i2 \rangle, \langle r, 0^i3 \rangle, \langle s, 0^i12 \rangle, \langle r, 0^i13 \rangle\}.$$

Suppose that  $ut_0^{\mathbb{S}}$  is realized by  $f : \Sigma^{\omega} \times \Sigma^{\omega} \rightarrow \Sigma^*$ . If  $\delta^u(p) = p_e^{x_i}$  and  $\delta^u(q) = p_e^{y_i}$ , then

$$f(p, q) \in \begin{cases} S_i & \text{if } i \in B \\ T_i & \text{if } i \in C. \end{cases} \quad (1)$$

$\forall i \in \mathbb{N}$  we define  $p_i = \iota(0^i1)\iota(0^i1)\dots$ , and  $q_i = \iota(0^i2)\iota(0^i2)\dots$ , where  $p_i, q_i \in \Sigma^{\omega}$ . Let  $F = \{f : \Sigma^{\omega} \times \Sigma^{\omega} \rightarrow \Sigma^* \text{ such that } f(p_i, q_i) \text{ exists for all } i \in \mathbb{N}\}$ . Consider  $f \in F$ . Then,  $f' : i \rightarrow f(p_i, q_i)$  is computable such that  $A \subseteq \text{dom}(f')$ . Since  $F$  is countable, there is a bijective function  $g : E \rightarrow F$  for some  $E \subseteq \mathbb{N}$  such that

$i \in \text{dom}(g'_i) \setminus A$  for all  $i \in E$ . Then,  $A \cap E = \emptyset$ . Let

$$B = \{i \in E : g_i(p_i, q_i) \notin S_i\}, C = \{i \in E : g_i(p_i, q_i) \in S_i\}, \quad (2)$$

and  $D = \mathbb{N} \setminus (A \cup B \cup C)$ . Since  $A \cap E = \emptyset$ ,  $E = B \cup C$  and  $B \cap C = \emptyset$ ,  $\{A, B, C, D\}$  is a partition of  $\mathbb{N}$ .

Suppose some computable function  $f$  realizes  $ut_0^s$ . Since  $\delta^u(p_i) = p_e^{x_i}$  and  $\delta^u(q_i) = p_e^{y_i}$  for all  $i \in A$ ,  $f(p_i, q_i)$  exists for all  $i \in A$ , hence  $f = g_i$  for some  $i \in E$ . Since  $g_i$  realizes  $ut_0^s$ ,  $g_i(p_i, q_i) \in S_i \Leftrightarrow i \in B$  by (3.1). On the other hand,  $g_i(p_i, q_i) \in S_i \Leftrightarrow i \notin B$  by 3.2. Thus, the space is not  $CuT_0$ .

**Example 3.14.** Let  $X = \{x\}$ ,  $E = \{e_1, e_2\}$  be a set of parameters, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  where  $\tau = \{\tilde{X}, \tilde{\emptyset}, \{(e_1, \{x\}), (e_2, \emptyset)\}\}$  which is generated by the following basis:

$$\nu(01) = \{(e_1, \{x\}), (e_2, \emptyset)\},$$

$$\nu(001) = \tilde{X},$$

Where  $\beta = \text{range}(\nu)$ . Thus,  $(X, \tau, E, \nu, \beta)$  is a computable STS and it is  $CuT_0$  but not  $CuT_1$ .

**Example 3.15.** Let  $A \subseteq \mathbb{N}$  be a c.e. set with non-c.e. complement. Define a notation  $\nu$  by

$$\nu(0^i 1) = \{p_e^{x_i}\}, \nu(0^i 2) = \{p_e^{x_i}\} \text{ for } i \in A,$$

$$\nu(0^i 1) = \{p_e^{x_i}\}, \nu(0^i 2) = \{p_e^{y_i}\} \text{ for } i \notin A,$$

for all  $i \in \mathbb{N}$ . Then,  $\nu$  is a notation of a base  $\beta$  of a STS on a subset  $X \subseteq \mathbb{N}$  w.r.t. a parameter set  $E = \{e\}$  such that  $(X, \tau, E, \beta, \nu)$  is a computable STS.

This space is  $CuT_2'$  as we have a c.e. set  $H = \{(0^i m, 0^i n) : i, j \in \mathbb{N}, m, n \in \{1, 2\}\}$  that satisfies  $CuT_2'$ . Let  $H$  be the c.e. set for  $SCuT_2$ . Thus, by the two conditions of  $SCuT_2$

$$i \notin A \Rightarrow (0^i 1, 0^i 2) \in H,$$

$$i \in A \Rightarrow (0^i 1, 0^i 2) \notin H,$$

since  $H$  is c.e., the complement of  $A$  must be c.e., which is a contradiction.

In the figure below, we summarize the all implications of the computable u-soft separation axioms. Those implications based on what we investigated above and the non implications are based on the counterexamples introduced

in this section above. These implications are actually the same as those of the classical computable separations axioms corresponding to the ones defined in the computable soft setting.




Figure 1: Relations between computable u-soft separation axioms

From the equivalences in figure 1, we can see that we have exactly four different notions of computable u-soft separation axioms.

In the next section, we will define computable p-soft separation axioms as the computable versions to those defined in [6]. Then, we define more variations of computable p-soft separations axioms and investigate the relations between them.

## 4 Computable p-soft separation axioms

In this section, we define the computable versions of partial soft separation axioms defined in [6] and then introduce some of the notions corresponding to those defined for computable u-soft separation axioms.

We define first  $\delta^p$  names for  $x_E \subseteq \tilde{X}$  in a computable STS  $(X, \tau, E, \beta, \nu)$ , where a  $\delta^p$  name of  $x_E \subseteq \tilde{X}$  contains all soft basic open sets that contain  $p_{e_i}^x$  for all  $e_i \in E$  where  $E$  is the parameter set associated with the given STS.

We will define also p-soft separation axioms based on  $x_E \subseteq \tilde{X}$  and then compare those separation axioms to the u-soft separation axioms defined in the previous section.

**Definition 4.1.** Let  $E$  be a finite set of parameters,  $\sigma$  be a function defined as follows  $\sigma : \subseteq \Sigma^* \setminus \{0\} \rightarrow E$  where  $\sigma(s_i) = e_i$ . Now,  $\delta^p(p) = x_E$  where  $p = s_i t(w_j) \dots$ , and  $p_{e_i}^x \in \nu(w_j)$ . In other words,  $p$  is a list of all soft basic open sets that contain  $p_{e_i}^x$  for all  $e_i \in E$ , and  $s_i$  tells us what parameter of  $E$  is represented.

Now, we define the p-soft separation axioms.

**Definition 4.2.** A computable STS  $(X, \tau, E, \beta, \nu)$  is

- computable p-soft  $T_0$  ( $CpT_0$ , for short) if  $(X, \tau, E)$  is a u-soft  $T_0$  and the multifunction  $pt_0$  is  $(\delta^p, \delta^p, \theta)$ -computable where  $pt_0$  maps every  $x_E, y_E \subseteq \tilde{X}$  such that  $x_E \neq y_E$  to some  $U_E \in \tau$  such that

$$(x \in U_E \text{ and } y \notin U_E) \text{ or } (x \notin U_E \text{ and } y \in U_E).$$

- computable p-soft  $T_1$  ( $CpT_1$ , for short) if  $(X, \tau, E)$  is a u-soft  $T_0$  and the multifunction  $pt_1$  is  $(\delta^p, \delta^p, \theta)$ -computable where  $pt_1$  maps every  $x_E, y_E \subseteq \tilde{X}$  such that  $x_E \neq y_E$  to some  $U_E \in \tau$  such that

$$(x \in U_E \text{ and } y \notin U_E).$$

- computable p-soft  $T_2$  ( $CpT_2$ , for short) if  $(X, \tau, E)$  is a u-soft  $T_0$  and the multifunction  $pt_2$  is  $(\delta^p, \delta^p, \theta)$ -computable where  $pt_2$  maps every  $x_E, y_E \subseteq \tilde{X}$  such that  $x_E \neq y_E$  to some  $U_E, V_E \in \tau$  such that

$$(x \in U_E \text{ and } y \in V_E, \text{ and } U_E \cap V_E = \tilde{\emptyset}).$$

We can see that  $CpT_i \Rightarrow CpT_{i-1}$  for  $i \in \{1, 2\}$ .

Based on the above definitions, we can see that the following implications hold,

$$CpT_2 \Rightarrow CpT_1 \Rightarrow CpT_0$$

The converses of the above implications are not true in general as shown from the following examples.

**Example 4.3.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  be a set of parameters and  $\tau$  is a STS defined on  $X$  w.r.t.  $E$  generated by the following basis,

$$\nu(01) = \{(e_1, \{x\}), (e_2, \emptyset)\}, \nu(001) = \{(e_1, \emptyset), (e_2, \{x\})\}, \nu(0001) = \tilde{X},$$

and  $\beta = \text{range}(\nu)$ . Thus,  $(X, \tau, E, \beta, \nu)$  is a computable STS and it is  $CpT_0$  since there is a machine  $M$  that realizes  $CpT_0$  where  $M$  on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$ , prints  $\iota(01)\iota(001)$ . The space is not  $CpT_1$  as it is not even  $pT_1$ .

**Example 4.4.** Let  $X = \{a_i : i \in \mathbb{N}\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  generated by the following basis notation,

$$\nu(o^i 1^j) = \{(e_1, G_i), (e_2, F_j)\},$$

where  $i$  and  $j$  are the canonical indices of  $G_i$  and  $F_j$ , respectively. We define the intersection of finitely many basic open sets by  $\nu(0^i 1^j) \cap \nu(0^k 1^l) = \nu(0^m 1^n)$ , where  $m$  is the canonical index of  $G_i^c \cap G_k^c$  and  $n$  is the canonical index of  $F_j \cap F_l$ . Thus, the space is computable STS. The space is  $CpT_1$  as there is a machine  $M$  that on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$ , searches for  $\iota(0^i 1^j)$  and  $\iota(0^m 1^n)$  such that  $\iota(0^i 1^j) \ll p$  and  $\iota(0^m 1^n) \ll q$ , and  $j$  and  $n$  are canonical indices of singletons of  $X$ , and  $i, m \in \mathbb{N}$ . If the search is successful, it prints  $\angle 0^n 1^j, 0^j 1^n \rangle$ . Hence, machine  $M$  realizes  $CpT_1$ . However, the space is not  $CpT_2$  as it is not  $pT_2$ .

Now, we give some more p-soft separation axioms and investigate the relations between them.

**Definition 4.5.** A computable STS  $(X, \tau, E, \beta, \nu)$  is:

$WCpT_0$ : if there is a c.e. set  $H \subseteq \text{dom}(\nu^{fs}) \times \text{dom}(\nu^{fs})$  such that

$$1. (\forall x_E \neq y_E)(\exists (u, v) \in H)(x \in \cup \nu^{fs}(u) \text{ and } y \in \cup \nu^{fs}(v)),$$

$$2. (\forall (u, v) \in H):$$

$$(\cup \nu^{fs}(u) \cap \cup \nu^{fs}(v) = \tilde{\emptyset}),$$

$$\vee ((\exists x_E) \cup \nu^{fs}(u) = x_E \subseteq \cup \nu^{fs}(v)),$$

$$\vee ((\exists y_E) \cup \nu^{fs}(v) = y_E \subseteq \cup \nu^{fs}(u)).$$

$SCpT_0$ : if the multifunction  $pt_0^s$  is  $(\delta^p, \delta^p, [\nu_N, \theta])$ -computable where  $pt_0^s$  maps every  $x_E, y_E \subseteq \tilde{X}$  such that  $(x_E \neq y_E)$  to some  $(k, U_E) \in \mathbb{N} \times \tau$  such that

$$(k = 1, x \in U_E \text{ and } y \notin U_E) \vee (k = 2, y \in U_E \text{ and } x \notin U_E).$$

$CpT'_0$ : if there is a c.e. set  $H \subseteq \text{dom}(\nu_N) \times \text{dom}(\nu^{fs}) \times \text{dom}(\nu^{fs})$  such that

$$1. (\forall x_E \neq y_E)(\exists (w, u, v) \in H)(x \in \cup \nu^{fs}(u) \text{ and } y \in \cup \nu^{fs}(v)),$$

2.  $(\forall(w, u, v) \in H)$ :

$$(\cup \nu^{fs}(u) \cap \cup \nu^{fs}(v) = \tilde{\emptyset}),$$

$$\vee(\nu_N(w) = 1(\exists x_E) \cup \nu^{fs}(u) = x_E \subseteq \cup \nu^{fs}(v)),$$

$$\vee(\nu_N(w) = 2(\exists y_E) \cup \nu^{fs}(v) = y_E \subseteq \cup \nu^{fs}(u)).$$

$CpT_1'$ : if there is a c.e. set  $H \subseteq \text{dom}(\nu^{fs}) \times \text{dom}(\nu^{fs})$  such that

1.  $(\forall x_E \neq y_E)(\exists(u, v) \in H)(x \in \cup \nu^{fs}(u) \text{ and } y \in \cup \nu^{fs}(v))$ ,

2.  $(\forall(u, v) \in H)$ :

$$(\cup \nu^{fs}(u) \cap \cup \nu^{fs}(v) = \tilde{\emptyset}),$$

$$\vee((\exists x_E) \cup \nu^{fs}(u) = x_E \subseteq \cup \nu^{fs}(v)).$$

$CpT_2'$ : if there is a c.e. set  $H \subseteq \text{dom}(\nu^{fs}) \times \text{dom}(\nu^{fs})$  such that

1.  $(\forall x_E \neq y_E)(\exists(u, v) \in H)(x \in \cup \nu^{fs}(u) \text{ and } y \in \cup \nu^{fs}(v))$ ,

2.  $(\forall(u, v) \in H)$ :

$$(\cup \nu^{fs}(u) \cap \cup \nu^{fs}(v) = \tilde{\emptyset}),$$

$$\vee((\exists x_E) \cup \nu^{fs}(u) = x_E = \cup \nu^{fs}(v)).$$

$SCP_{T_2}$ : if there is a c.e. set  $H \subseteq \text{dom}(\nu^{fs}) \times \text{dom}(\nu^{fs})$  such that

1.  $(\forall x_E \neq y_E)(\exists(u, v) \in H)(x \in \cup \nu^{fs}(u) \text{ and } y \in \cup \nu^{fs}(v))$ ,

2.  $(\forall(u, v) \in H)$ :

$$(\cup \nu^{fs}(u) \cap \cup \nu^{fs}(v) = \tilde{\emptyset}).$$

**Proposition 4.6.** *Let  $\overline{CpT}_i$  and  $\overline{SCP}_{T_0}$  be the conditions obtained from  $CpT_i$  and  $SCP_{T_0}$ , respectively, by replacing  $\theta$  by  $\cup \nu^{fs}$ . Then,  $\overline{CpT}_i \Leftrightarrow CpT_i$  for  $i \in \{0, 1, 2\}$ , and  $\overline{SCP}_{T_0} \Leftrightarrow SCP_{T_0}$ , when the parameter set is finite.*

*Proof.*  $\overline{CpT}_i \Rightarrow CpT_i$ : since  $\cup \nu^{fs} \leq \theta$ .

$CpT_i \Rightarrow \overline{CpT}_i$ : There is a machine  $M$  that on input  $(p, q) \in \text{dom}(\delta^p) \times \text{dom}(\theta)$  where  $\delta^p(p) \in \theta(q)$  searches for  $u_1, \dots, u_n$  where  $u_i \ll p_i$  and  $u_i \ll q$  for all  $i$  where  $p_i$  is a  $\delta^u$  name obtained from  $p$ . Then, machine  $M$  prints  $u$  if the search is successful where  $u = \iota(u_1)\iota(u_2)\dots$  and diverges, otherwise. Following the same argument, we can prove  $\overline{SCP}_{T_0} \Leftrightarrow SCP_{T_0}$ , which completes the proof.  $\square$

We now introduce some implications between the p-soft spaces defined above.

**Proposition 4.7.**  $CpT_0 \Leftrightarrow SCpT_0 \Leftarrow CpT'_0 \Rightarrow WCpT_0$ .

*Proof.*  $SCpT_0 \Rightarrow CpT_0$ : Obvious.

$CpT_0 \Rightarrow SCpT_0$ : There is a machine  $M$  that on input  $(p, q) \in \text{dom}(\delta^p) \times \text{dom}(\delta^p)$  searches for  $u_1, \dots, u_n \in \text{dom}(\nu)$  such that  $u_i \ll r$  for all  $i \in \{1, \dots, n\}$  where  $pt_0(p, q) = r$ , and outputs  $\langle 1, r \rangle$  if for all  $i$   $u_i \ll p_i$  where  $p_i$  is a  $\delta^u$ -name obtained from  $p$ , and outputs  $\langle 2, r \rangle$  if for all  $i$   $u_i \ll q_i$  where  $q_i$  is a  $\delta^u$ -name obtained from  $q$ . We can see easily the  $M$  realizes  $pt_0^s$ , which completes the proof.

$CpT'_0 \Rightarrow SCpT_0$ : There is a machine  $M$  that on input  $(p, q) \in \text{dom}(\delta^p) \times \text{dom}(\delta^p)$  searches for  $(w, r, s) \in H$ -The c.e. set of  $CpT'_0$ - such that there are for all  $i \in \{1, \dots, n\}$   $u_i, v_i \in \text{dom}(\nu)$  where  $u_i \ll r$ ,  $v_i \ll s$  and  $u_i \ll p_i, v_i \ll q_i$  where  $p_i, q_i$  are  $\delta^u$ -names obtained from  $p, q$ , respectively. Then, machine  $M$  prints  $\langle w, r \rangle$  if  $\nu_N(w) = 1$  and  $\langle w, s \rangle$ , otherwise. Thus  $M$  realizes  $pt_0^s$ , which completes the proof.

$CpT'_0 \Rightarrow SCpT_0$ : Obvious.  $\square$

We now show that the second and third implications are not reversed in general as shown from the next two examples.

**Example 4.8.** Let  $X = \{a_i, b_i : i \in \mathbb{N}\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  generated by the following basis where  $A$  is a non c.e. set,

	$0^i 11$	$0^i 12$	$0^i 51$	$0^i 52$	$0^i 5211$
$i \in A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$\tilde{\emptyset}$
$i \notin A$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$p_{e_1}^{x_i}$	$p_{e_1}^{y_i} \cup p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$

The finite intersections are all empty except for  $\nu(0^i 11) \cap \nu(0^i 52) = \nu(0^i 5211)$ . Thus, the space  $(X, \tau, E, \beta, \nu)$  is computable STS. Let  $H$  be the c.e. set of  $WCpT_0$ , then

$$i \in A \Rightarrow (\iota(0^i 11)\iota(0^i 12), \iota(0^i 51)\iota(0^i 52)) \in H,$$

$$i \notin A \Rightarrow (\iota(0^i 11)\iota(0^i 12), \iota(0^i 51)\iota(0^i 52)) \notin H.$$

Thus,  $A$  must be an r.e. set which is a contradiction. Hence, the space is not  $WCpT_0$  and then not  $CpT'_0$ , however, It is  $CpT_0$  as there is a machine  $M$  that

realizes  $pt_0$  where  $M$  on  $(p, q)$  prints  $\iota(0i11)\iota(0^i12)$ .

**Proposition 4.9.** *There is a computable STS that is  $WCpT_0$  but not  $CpT_0$ .*

*Proof.* Follows immediately from the following example.  $\square$

**Example 4.10.** *Let  $A \subseteq \mathbb{N}$  be some non c.e. set. Let  $X = \{x_i, y_i\}$ ,  $E = \{e_1, e_2\}$  be a parameter set and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  generated by the following basis given in the table below.*

	$0^i11$	$0^i12$	$0^i21$	$0^i22$	$0^i31$	$0^i32$
$i \in A \cup D$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$
$i \in B$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{x_i} \cup p_{e_1}^{y_i}$	$p_{e_2}^{x_i} \cup p_{e_2}^{y_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$
$i \in C$	$p_{e_1}^{x_i} \cup p_{e_1}^{y_i}$	$p_{e_2}^{x_i} \cup p_{e_2}^{y_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$

We define  $\{A, B, C, D\}$  to be a partition of  $\mathbb{N}$ . We define the intersection of soft basic open sets as follows,  $\nu(0^i kl) \cap \nu(0^i mn) = \nu(0^i klmn)$  for  $k \neq m \vee l \neq n$ . Therefore,  $(X, \tau, E, \beta, \nu)$  is a computable STS. We can see that the space is  $WCpT_0$  as we can have a c.e. set  $H = \{\langle \iota(0^i r1)\iota(0^i r2), \iota(0^j s1)\iota(0^j s2) \rangle : i, j \in \mathbb{N}; r, s \in \{1, 2\}; (i \neq j \vee r \neq s)\}$  that satisfies the two conditions of  $WCpT_0$ . Now, we define  $B$  and  $C$  in a way that makes the space not  $SCpT_0$ . Let  $w_1, w_2 \in \Sigma^8$  such that  $\nu_N(w_1) = 1$  and  $\nu_N(w_2) = 2$ , and W.L.O.G. we assume that  $\nu_N$  is injective. For  $i \in \mathbb{N}$  let  $S_i = \{\langle w_1, \iota(0^i 11)\iota(0^i 12) \rangle, \langle w_1, \iota(0^i 11)\iota(0^i 1222) \rangle, \langle w_1, \iota(0^i 1121)\iota(0^i 12) \rangle, \langle w_1, \iota(0^i 1121)\iota(0^i 1222) \rangle, \langle w_2, \iota(0^i 31)\iota(0^i 32) \rangle, \langle w_2, \iota(0^i 31)\iota(0^i 2232) \rangle, \langle w_2, \iota(0^i 2131)\iota(0^i 32) \rangle, \langle w_2, \iota(0^i 2131)\iota(0^i 2232) \rangle\}$ ,  $T_i = \{\langle w_1, \iota(0^i 31)\iota(0^i 32) \rangle, \langle w_1, \iota(0^i 31)\iota(0^i 1232) \rangle, \langle w_1, \iota(0^i 1131)\iota(0^i 32) \rangle, \langle w_1, \iota(0^i 1131)\iota(0^i 1232) \rangle, \langle w_2, \iota(0^i 21)\iota(0^i 22) \rangle, \langle w_2, \iota(0^i 21)\iota(0^i 1222) \rangle, \langle w_2, \iota(0^i 1121)\iota(0^i 22) \rangle, \langle w_2, \iota(0^i 1121)\iota(0^i 1222) \rangle\}$

Suppose the function  $f : \subseteq \Sigma^\omega \times \Sigma^\omega \rightarrow \Sigma^*$  realizes  $pt_0^s$ . If  $\delta^p(p) = a_{E_i}$  and  $\delta^p(q) = b_{E_i}$ , then

$$f(p, q) \in \begin{cases} S_i & \text{if } i \in B \\ T_i & \text{if } i \in C. \end{cases} \quad (3)$$

$\forall i \in \mathbb{N}$  we define  $p_i = \iota(0^i 11)\iota(0^i 12)\iota(0^i 11)\iota(0^i 12) \cdots$ ,  
and  $q_i = \iota(0^i 21)\iota(0^i 22)\iota(0^i 21)\iota(0^i 22) \cdots$ , where  $p_i, q_i \in \Sigma^\omega$ . Let  $F = \{f : \Sigma^\omega \times \Sigma^\omega \rightarrow \Sigma^* \text{ such that } f \text{ is computable and } f(p_i, q_i) \text{ exists for all } i \in A\}$ .  
Consider  $f \in F$ . Then,  $f' : i \rightarrow f(p_i, q_i)$  is computable such that  $A \subseteq \text{dom}(f')$ .  
Since  $F$  is countable, there is a bijective function  $g : E \rightarrow F$  for some  $E \subseteq \mathbb{N}$   
such that  $i \in \text{dom}(g'_i) \setminus A$  for all  $i \in E$ . Then,  $A \cap E = \emptyset$ . Let

$$B = \{i \in E : g_i(p_i, q_i) \notin S_i\}, C = \{i \in E : g_i(p_i, q_i) \in S_i\}, \quad (4)$$

and  $D = \mathbb{N} \setminus (A \cup B \cup C)$ . Since  $A \cap E = \emptyset$ ,  $E = B \cup C$  and  $B \cap C = \emptyset$ ,  
 $\{A, B, C, D\}$  is a partition of  $\mathbb{N}$ .

Suppose some computable function  $f$  realizes  $pt_0^s$ . Since  $\delta^p(p_i) = x_{E_i}$  and  
 $\delta^p(q_i) = y_{E_i}$  for all  $i \in A$ ,  $f(p_i, q_i)$  exists for all  $i \in A$ , hence  $f = g_i$  for some  
 $i \in E$ . Since  $g_i$  realizes  $pt_0^s$ ,  $g_i(p_i, q_i) \in S_i \Leftrightarrow i \in B$  by (3.3). On the other hand,  
 $g_i(p_i, q_i) \in S_i \Leftrightarrow i \notin B$  by 3.4. Thus, the space is not  $CpT_0$ .

**Proposition 4.11.**  $SCpT_2 \Rightarrow CpT_2 \Leftarrow CpT'_2 \Leftrightarrow CpT'_1 \Rightarrow CpT_1$ .

*Proof.*  $SCpT_2 \Rightarrow CpT_2$ : There is a machine  $M$  that on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$   
searches for  $(r, s) \in H$  such that  $\forall i$  there are  $u_i, v_i \in \text{dom}(\nu)$  for  $i \in \{0, 1, \dots, n\}$   
where  $u_i \ll r$ ,  $v_i \ll s$  and  $u_i \ll p_i$ ,  $v_i \ll q_i$ , where  $p_i, q_i$  are  $\delta^u$ -names obtained  
from  $p, q$ , respectively. Machine  $M$  prints  $\langle r, s \rangle$  if the search successful and  
diverges, otherwise.

Thus, let  $\delta^p(p) = x_E \neq y_E = \delta^p(q)$ . When we apply  $M$  on  $(p, q)$ , the machine  
searches for  $(r, s) \in H$  as described above and the search must be successful  
since by definition of  $H$  there must exist  $(r, s) \in H$  such that  $x \in \cup \nu^{fs}(r)$  and  
 $y \in \cup \nu^{fs}(s)$  and  $\cup \nu^{fs}(r) \cap \cup \nu^{fs}(s) = \tilde{\emptyset}$ , and thus,  $\forall i \in \{1, 2, \dots, n\}$  there exists  
 $u_i \ll r$ ,  $v_i \ll s$  such that  $p_{e_i}^x \in \nu(u_i)$  and  $p_{e_i}^y \in \nu(v_i)$ . Therefore, the space is  
 $CpT_2$ .

$CpT'_2 \Rightarrow CpT_2$ : There is a machine  $M$  on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$  searches for  
 $(r, s) \in H$  such that  $\forall i \in \{1, 2, \dots, n\}$  there are  $u_i, v_i \in \text{dom}(\nu)$  and  $u_i \ll p_i$ ,  
 $v_i \ll q_i$ . The machine prints  $\langle r, s \rangle$  if the search is successful and diverges,  
otherwise. Thus, machine  $M$  realizes  $pt_2$ .

$CpT'_2 \Rightarrow CpT'_1$ : Obvious.

$CpT'_1 \Rightarrow CpT'_2$ : We define the c.e. set of  $CpT'_2$  to be  $H_2 = \{(\bar{r}, \bar{s}) : u_i \ll \bar{r} \Rightarrow$   
 $u_i \ll g(r, s'), v_i \ll \bar{s} \Rightarrow g(r', s) \text{ for some } (r, s), (r', s') \in H\}$  where  $H$  is the c.e.  
set of  $CpT'_1$ .

We check now the two conditions of  $H_2$ . Let  $x_E \neq y_E$ . There are

$(r, s), (r', s') \in H$  such that  $x \in \cup \nu^{fs}(r) \cap \cup \nu^{fs}(s')$ ,  $y \in \cup \nu^{fs}(s) \cap \cup \nu^{fs}(r')$ . Then,  $x \in \theta(r'') = \cup \nu^{fs}(r) \cap \cup \nu^{fs}(s')$ ,  $y \in \theta(s'') = \cup \nu^{fs}(s) \cap \cup \nu^{fs}(r')$  and hence  $\forall i \in \{1, 2, \dots, n\}$  there are  $u_i \ll r''$  and  $v_i \ll s''$  where  $p_{e_i}^x \in \nu(u_i)$  and  $p_{e_i}^y$ . Thus, there is  $(\bar{r}, \bar{s})$  where  $\bar{r} = \iota(u_1) \dots \iota(u_n)$ ,  $\bar{s} = \iota(v_1) \dots \iota(v_n)$  and  $x \in \cup \nu^{fs}(\bar{r})$ ,  $y \in \cup \nu^{fs}(\bar{s})$ .

Now, we prove the second condition of  $H_2$ . Suppose  $(\bar{r}, \bar{s}) \in H_2$  and  $\cup \nu^{fs}(\bar{r}) \cap \cup \nu^{fs}(\bar{s}) \neq \tilde{\emptyset}$ . Thus, there are  $(r, s), (r', s') \in H$  such that  $\cup \nu^{fs}(\bar{r}) \subseteq \cup \nu^{fs}(r) \cap \cup \nu^{fs}(s')$ , and  $\cup \nu^{fs}(\bar{s}) \subseteq \cup \nu^{fs}(r') \cap \cup \nu^{fs}(s)$ , and then  $\cup \nu^{fs}(r) \cap \cup \nu^{fs}(s) \neq \tilde{\emptyset}$ , and  $\cup \nu^{fs}(r') \cap \cup \nu^{fs}(s') \neq \tilde{\emptyset}$ . Hence, there are  $x_E$  and  $y_E$  such that  $\cup \nu^{fs}(r) = x_E \subseteq \cup \nu^{fs}(s)$ , and  $\cup \nu^{fs}(r') = y_E \subseteq \cup \nu^{fs}(s')$ . Therefore,  $\cup \nu^{fs}(\bar{r}) \subseteq x_E$  and  $\cup \nu^{fs}(\bar{s}) \subseteq y_E$ , which means that  $x_E = y_E$ . Thus, the second condition of  $H_2$  is satisfied.

$CpT'_1 \Rightarrow CpT_1$ : This is a special case of  $CpT'_0 \Rightarrow SCpT_0$ , which completes the proof.  $\square$

We introduce now counterexamples to show that the implications of the previous proposition are not reversed in general.

The next example shows a space which is  $CpT_2$  but not  $SCpT_2$ .

**Remark 4.12.**  $CpT'_2 \Rightarrow CpT'_0$

*Proof.* Straightforward.  $\square$

**Example 4.13.** Let  $A \subseteq \mathbb{N}$  be a c.e. set with non c.e. complement. We define a notation of a basis of a topology  $\tau$  on a subset  $X \subseteq \mathbb{N}$  as follows,

	$0^i 11$	$0^i 12$	$0^i 21$	$0^i 22$
$i \in A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$
$i \notin A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$

Thus,  $(X, \tau, E, \beta, \nu)$  is a computable STS. The space is  $CpT'_2$  as we have a c.e. set that satisfies the two conditions of it, namely,

$$H = \{(\iota(0^i k l) \iota(0^i m n)), (\iota(0^j k' l') \iota(0^j m' n')) : i, j \in \mathbb{N}; k, l, m, n, k', l', m', n' \in \{1, 2\}; ((k = m \text{ and } k' = m') \text{ and } (k \neq l \vee m \neq n) \text{ and } (k \neq l \vee m' \neq n'))\}.$$

Now, we show that the space is not  $SCpT_2$ . Let  $H'$  be the c.e. set of  $SCpT_2$ . then by the first condition of  $H'$ ,

$$i \notin A \Rightarrow (u, v) \in H' \text{ where } 0^i 11, 0^i 12 \ll u \text{ and } 0^i 21, 0^i 22 \ll v,$$

and by the second condition of  $H'$ ,

$$i \in A \Rightarrow (u, v) \notin H' \text{ where } 0^i 11, 0^i 12 \ll u \text{ and } 0^i 21, 0^i 22 \ll v.$$

Since  $H'$  c.e., the complement of  $A$  must be c.e. which a contradiction.

The next example shows that there is a space which is  $CpT_2$  but not  $CpT'_1$ .

**Example 4.14.** Let  $A \subseteq \mathbb{N}$  be a non c.e. set,  $E = \{e_1, e_2\}$  be a parameter set and  $X = \{x_i, y_i : i \in \mathbb{N}\}$  be a set on which a STS  $\tau$  is defined where  $\tau$  is generated by the following basis which is given by the the following notation,

	$0^i 11$	$0^i 12$	$0^i 13$	$0^i 14$	$0^i 1112$	$0^i 1113$	$0^i 1114$
$i \in A$	$x_{E_i}$	$y_{E_i}$	$p_{e_1}^{x_i}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$p_{e_1}^{x_i}$	$\tilde{\emptyset}$
$i \notin A$	$x_{E_i} \cup p_{e_1}^{y_i}$	$x_{E_i} \cup p_{e_2}^{y_i}$	$\tilde{\emptyset}$	$p_{e_2}^{y_i}$	$x_{E_i}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$

	$0^i 1213$	$0^i 1214$	$0^i 1314$	$0^i 61$	$0^i 6111$	$0^i 6112$	$0^i 6113$	$0^i 6114$
$i \in A$	$\tilde{\emptyset}$							
$i \notin A$	$\tilde{\emptyset}$	$p_{e_2}^{y_i}$	$\tilde{\emptyset}$	$p_{e_1}^{y_i}$	$p_{e_1}^{y_i}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$

Thus,  $(X, \tau, E, \beta, \nu)$  is a computable STS. The space is  $CpT_2$  as there is a machine  $M$  that on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$  searches for  $0^i 13$  and  $0^i 14$  where on of the following cases hold:

1.  $0^i 13 \ll p$  and:

- (a)  $0^i 12 \ll q$ , the machine prints  $\langle 0^i 11, 0^i 12 \rangle$ ,
- (b)  $0^j 12 \vee 0^j 11 \ll q$  for some  $j \neq i$ , the machine prints  $\langle 0^j 11, \iota(0^j 11 \iota(0^j 12)) \rangle$

2.  $0^i 13 \ll q$  and:

- (a)  $0^i 12 \ll p$ , the machine prints  $\langle 0^i 12, 0^i 11 \rangle$ ,
- (b)  $0^j 12 \vee 0^j 11 \ll p$  for some  $j \neq i$ , the machine prints  $\langle \iota(0^j 11 \iota(0^j 12)), 0^j 11 \rangle$

3.  $0^i 14 \ll p$  and:

(a)  $0^i 1112 \ll q$ , the machine prints  $\langle \iota(0^i 14)\iota(0^i 61), 0^i 1112 \rangle$

(b)  $0^j 12 \vee 0^j 11 \ll q$  for  $j \neq i$ , the machine prints  $\langle \iota(0^i 11)\iota(0^i 12), 0^i 1112 \rangle$

4.  $0^i 14 \ll q$  and:

(a)  $0^i 1112 \ll p$ , the machine prints  $\langle 0^i 1112, \iota(0^i 14)\iota(0^i 61) \rangle$

(b)  $0^j 12 \vee 0^j 11 \ll p$  for  $j \neq i$ , the machine prints  $\langle 0^i 1112, \iota(0^i 11)\iota(0^i 12) \rangle$

Hence,  $M$  realizes  $CpT_2$ . Now, we prove that the space is not  $CpT'_1$ . Let  $H$  be the c.e. set of  $CpT'_1$ , then

$$i \in A \Rightarrow (u, v) \in H \text{ where } 0^i 11 \ll u, 0^i 12 \ll v,$$

$$i \notin A \Rightarrow (u, v) \notin H \text{ where } 0^i 11 \ll u, 0^i 12 \ll v.$$

Since  $H$  is a c.e. set,  $A$  must be a c.e. set which is a contradiction. Therefore, the space is not  $CpT'_1$ .

In the following figure, we represent all implications between the computable p-soft separation axioms we defined so far. The implications are based on the results that we got in this section and the non implications come from the counterexamples that we introduced above.




Figure 2: Relations between computable p-soft separation axioms

We can see from fig 2 that we have exactly seven different notions of p-soft separation axioms compared to four different notions of u-soft separation axioms.

In the next section, we study the relation between computable u-soft separation axioms and computable p-soft separation axioms.

## 5 Relations between u-soft and p-soft separation axioms

In this section, we investigate how computable u-soft separation axioms are related to their counterparts computable p-soft separation axioms. We just consider the case when the set of parameters is finite.

At the end of this section, we will be able to compare the four different notions of u-soft separation axioms to the seven different notions of the p-soft separation axioms.

**Proposition 5.1.** *Computable u-soft  $T_i \Rightarrow$  computable p-soft  $T_i$ , for  $i = 1, 2$ .*

*Proof.* Case 1:  $i = 1$ . Assume u-soft  $T_1$ . Let  $\delta^p(p) = x_E \neq y_E = \delta^p(q)$ . There are  $n$  machines  $M_i$  such that machine  $M_i$  translates  $p$  into a  $\delta^u$ -name  $p_i$  for  $p_{e_i}^x$ . Similarly, there are  $n$  machines  $N_i$  where  $N_i$  translates  $q$  into a  $\delta^u$ -name for  $p_{e_i}^y$ . Now,  $\forall i \forall j$   $ut_1$  on input  $(p_i, q_i)$  outputs  $w_{i_j}$  where  $\nu(w_{i_j}) = U_{E_{i_j}} \in \beta$  and  $p_{e_i}^x \in U_{E_{i_j}}$  and  $p_{e_j}^y \notin U_{E_{i_j}}$ .  $\forall i$ , let  $w_i = \iota(w_{i_1}) \dots \iota(w_{i_n})$  and since  $\nu \leq \theta$  and the intersection of a finite set of open sets is  $(\theta^{fs}, \theta)$ -computable, there is a computable function  $f$  such that  $\bigcap \nu^{fs}(w_i) = \theta \circ f(w_i)$ . Thus,  $\forall i \forall j, p_{e_i}^x \in \theta(r_i)$  and  $p_{e_j}^y \notin \theta(r_i)$  where  $r_i = f(w_i)$ . Also, since the union of a finite set of open sets is open, there is a computable function  $g$  such that  $\bigcup \theta^{fs}(\langle 1^n, r_1, \dots, r_n \rangle) = \theta \circ g(\langle 1^n, r_1, \dots, r_n \rangle)$  and hence  $x \in \theta(r)$  and  $y \notin \theta(r)$  where  $r = \langle 1^n, r_1, \dots, r_n \rangle$ . Therefore, the space is p-soft  $T_1$ .

Case 2:  $i = 2$ . Assume u-soft  $T_2$ . Let  $\delta^p(p) = x_E \neq y_E = \delta^p(q)$ . There are  $n$  machines  $M_i$  such that machine  $M_i$  translates  $p$  into a  $\delta^u$ -name  $p_i$  for  $p_{e_i}^x$ . Similarly, there are  $n$  machines  $N_i$  where  $N_i$  translates  $q$  into a  $\delta^u$ -name for  $p_{e_i}^y$ .  $\forall i \forall j$   $ut_2$  on input  $(p_i, q_j)$  outputs  $(u_{i_j}, v_{i_j})$  where  $\nu(u_{i_j}) = G_{E_{i_j}} \in \beta$  and  $\nu(v_{i_j}) = H_{E_{i_j}} \in \beta$  such that  $p_{e_i}^x \in G_{E_{i_j}}$  and  $p_{e_j}^y \in H_{E_{i_j}}$  and  $G_{E_{i_j}} \cap H_{E_{i_j}} = \tilde{\emptyset}$ .  $\forall i$ , let  $u_i = \iota(u_{i_1}) \dots \iota(u_{i_n})$  and  $v_i = \iota(v_{i_1}) \dots \iota(v_{i_n})$ . By functions  $f$  and  $g$  from case 1,  $\forall i$  we have  $f(u_i) = r_i$  and  $g(v_i) = s_i$  where  $p_{e_i}^x \in \theta(r_i)$  and  $y \in \theta(s_i)$  and  $\theta(r_i) \cap \theta(s_i) = \tilde{\emptyset}$ . Now, we use  $g$  and  $f$  again, where  $g(\langle 1^n, r_1, \dots, r_n \rangle) = r$  and  $f(\langle 1^n, s_1, \dots, s_n \rangle) = s$ . Thus,  $x \in \theta(r)$  and  $y \in \theta(s)$  and  $\theta(r) \cap \theta(s) = \tilde{\emptyset}$ . Therefore, the space is  $p$ -soft  $T_2$  which completes the proof.  $\square$

We give a counter example that the converse of the above implications is not true in general.

**Example 5.2.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  and generated by the following base  $\nu(01) = \{(e_1, \{x\}), (e_2, \{x\})\}$ ,  $\nu(001) = \{(e_1, \{y\}), (e_2, \{y\})\}$ .  $\nu(001) = \{(e_1, \{x\}), (e_2, \{y\})\}$ . The space is computable STS and it is  $CpT_1$  as we have a machine  $M$  on  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$  outputs  $\iota(01)$  if  $01 \ll p$  and outputs  $\iota(001)$  if  $01 \ll q$ . Thus,  $M$  realizes  $pt_1$  but the space is not  $CuT_1$  as it is not  $u$ -soft  $T_1$ . We can see also that this space is  $CpT_2$  but not  $CuT_2$ .

In the next example, we show that the above result does not hold when the set of parameters is infinite.

**Example 5.3.** Let  $X = \{a, b\}$ ,  $E = \{e_1, e_2, \dots\}$ . We partition  $\mathbb{N}$  into infinitely many infinite partitions  $\mathbb{N} = F_1 \cup F_2 \cup \dots$ . we define a STS on  $X$  with respect to  $E$  where its basic open sets are defined as follows, for each finite set  $G \subseteq \mathbb{N}$  we have  $\{p_{e_i}^a : i \in G\}$  and for each finite set  $G \subseteq \mathbb{N}$ ,  $n \in \mathbb{N}$  we have  $\{p_{e_n}^b\} \cup \{p_{e_i}^a : i \in F_n - G\}$ . Clearly, this space is  $u$ -soft  $T_2$  but it is not  $p$ -soft  $T_2$ . We effectivize this space by introducing a notation  $\nu$  for the set of basic open sets  $\beta$  as follows,  $\nu(0^k1) = G$  where  $k$  is the canonical index of  $G$ , and  $\nu(0^m10^n1) = \{p_{e_m}^b\} \cup \{p_{e_i}^a : i \in F_m - G\}$  where  $m$  is the index of  $F_m$  and  $n$  is the canonical index of  $G$ . We define the finite intersection of basic open sets as follows,

- $\nu(0^k1) \cap \nu(0^l1) = \nu(0^r1)$  where  $r$  is the canonical index of the intersection of two sets, the canonical index of the first set is  $k$  while the canonical index of the other one is  $l$ .
- $\nu(0^m10^n1) \cap \nu(0^r10^s1) = \tilde{\emptyset}$  for  $m \neq n$ .
- $\nu(0^m10^n1) \cap \nu(0^r10^s1) = \nu(0^m10^t1)$  for  $m = n$ , where  $t$  is the canonical index of the set resulting from the union of two sets whose canonical indices are  $s$  and  $n$ .
- $\nu(0^k1) \cap \nu(0^m10^n1) = \nu(0^s1)$  where  $s$  is the canonical index of  $H$  where  $H = G \cap F_m - I$  and  $k, n$  are the canonical indices of  $G, I$ , respectively, and  $m$  is the index of  $F_m$ .

Finite intersections can be obtained directly from the cases above. Thus, the space  $(X, \tau, E, \nu, \beta)$  is a computable STS.

Now, we show that the space is  $CuT_2$ . There is a machine  $M$  that on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$  scans the first words  $u, v$  of  $p, q$  respectively, until it finds one of the following,

- $u, v$  of the form  $0^i1$  for some  $i$ , then the machine scans through  $p, q$  until it finds  $0^r1 \ll p$  and  $0^s1 \ll q$  where  $r, s$  are canonical indices of singletons. Then, it outputs  $\langle 0^r1, 0^s1 \rangle$ .
- $u, v$  of the form  $0^i10^j1, 0^m10^n1$ , respectively. If  $i = m$ , then the machine scans through  $p, q$  until it finds  $0^r1 \ll p$  and  $0^s1 \ll q$  where  $r, s$  are canonical indices of singletons and outputs  $\langle 0^r1, 0^s1 \rangle$ . It outputs  $\langle u, v \rangle$ , otherwise.
- $u$  is of the form  $0^i1$  and  $v$  is of the form  $0^m10^n1$ , then  $M$  scans through  $p, q$  until it finds  $0^j1 \ll p$  and  $0^m10^j1 \ll q$  where  $j$  is the canonical index of a singleton. Then, it outputs  $\langle 0^j1, 0^m10^j1 \rangle$ .

Thus, machine  $M$  realizes  $ut_2$  and hence the space is  $CuT_2$ . However, the space is not even  $p$ -soft  $T_1$ .

The following two examples show that  $CuT_0$  and  $CpT_0$  are incomparable.

**Example 5.4.** Let  $X = \{x_i, y_i : i \in \mathbb{N}\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  and generated by the following base notation where  $A$  is a non c.e. set,

	$0^i11$	$0^i12$	$0^i21$	$0^i22$	$0^i31$	$0^i1131$	$0^i2131$	$0^i1112$	$0^i1231$
$i \in A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$	$\tilde{\emptyset}$
$i \notin A$	$x_{E_i}$	$x_{E_i}$	$y_{E_i}$	$\tilde{\emptyset}$	$p_{e_1}^{x_i} \cup p_{e_2}^{y_i}$	$p_{e_1}^{x_i}$	$p_{e_2}^{y_i}$	$x_{E_i}$	$p_{e_1}^{x_i}$

We extend names to the finite intersections as follows  $\nu(0^i mn) \cap \nu(0^i kl) = \nu(0^i mnkl)$  and the intersection of more than two basic open sets is empty except for  $\nu(0^i11) \cap \nu(0^i12) \cap \nu(0^i31) = \nu(0^i111231)$ . Thus, the space is computable STS. The space is  $CpT_2$ - and hence  $CpT_0$ - as we have the following c.e. set,

$$H = \{(\iota(0^i m1)\iota(0^i m2), \iota(0^i n1)\iota(0^i n2)) : m, n \in \{1, 2\}; i, j \in \mathbb{N}\}.$$

Assume now that the space is  $WCuT_0$ . Thus, there is a c.e. set  $H^i$  that satisfies

the two conditions of  $WCuT_0$  which means the following,

$$i \in A \Rightarrow (0^i 11, 0^i 12) \in H',$$

$$i \notin A \Rightarrow (0^i 11, 0^i 12) \notin H'.$$

Hence,  $A$  must be a c.e. set which is a contradiction. Therefore the space is not  $WCuT_0$  (Thus not  $CuT_0$  as well).

**Example 5.5.** Let  $X = \{x, y\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  and generated by the following base notation,

$$\begin{aligned} \nu(01) &= \{(e_1, \{x\}), (e_2, \emptyset)\}, \\ \nu(02) &= \{(e_1, \{y\}), (e_2, \emptyset)\}, \\ \nu(03) &= \{(e_1, \{x\}), (e_2, \{y\})\}, \\ \nu(04) &= \{(e_1, \{y\}), (e_2, \{x\})\}, \end{aligned}$$

We give names to the finite intersections of basic open sets as follows,  $\nu(0m) \cap \nu(0n) = \nu(0mn)$  for  $m, n \in \{1, \dots, 4\}$  and the intersection of any three basic open sets is empty.

Now, we show that this space is  $u$ -soft  $T_0$ . There is a machine  $M$  that on input  $(p, q) \in \Sigma^\omega \times \Sigma^\omega$ , scans  $p$  and  $q$  and prints  $\iota(u)$  whenever it scans first  $u \ll p$  or  $u \ll q$  such that  $u \in \{01, 02\}$  at any point of the computation. If  $M$  scans first  $0m \ll p$  or  $0m \ll q$  for  $m \in \{3, 4\}$ , then it prints the first word  $v$  of the other name if  $v \neq 0m$ , otherwise, it prints  $\iota(01)$  if  $m = 3$  and prints  $\iota(02)$  if  $m = 4$ .

Therefore, machine  $M$  realizes  $ut_0$  and hence the space is  $CuT_0$ . However, it is not  $CpT_0$  as it is not  $pT_0$ .

**Proposition 5.6.**  $SCuT_2 \Rightarrow SCpT_2$ .

*Proof.* Let  $H$  be the c.e. set of  $SCuT_2$ , and  $n$  be the number of parameters. Let  $H' \subseteq \Sigma^* \times \Sigma^*$  be the set of all pairs  $(u, v)$  of words for which there are some  $n$  such that  $u = \iota(u_1) \dots \iota(u_n)$  and  $v \ll \theta \circ f(\iota(v_1) \dots \iota(v_n))$ , where  $f$  is the computable function that computes the finite intersection of soft open set and  $u_1, \dots, u_2 \in \text{dom}(\nu)$  and  $v_1, \dots, v_n \in \text{dom}(\nu^{fs})$ , and  $\forall i (u_i \ll \bigcap \nu^{fs}(w_i) \text{ where } \nu^{fs}(w_i) = Pr_1(N) \text{ and } \nu^{fs}(v_i) = Pr_2(N) \text{ for some finite set } N \subseteq H$ .

Let  $x_E \neq y_E$ . Then,  $\forall p_{e_i}^x \in x_E \forall p_{e_j}^y \in y_E$  there are pairs  $(r_{i_1}, s_{i_1}), \dots, (r_{i_n}, s_{i_n}) \in H$  such that  $p_{e_i}^x \in \nu(r_{i_j})$  and  $p_{e_j}^y \in \nu(s_{i_j})$ , and  $\nu(r_{i_j}) \cap \nu(s_{i_j}) = \emptyset$ . Then,  $p_{e_i}^x \in \bigcap \nu^{fs}(w_i)$  where  $w_i = \iota(r_{i_1}) \dots \iota(r_{i_n})$  and

hence there is some  $u_i \ll \nu^{fs}(w_i)$  where  $p_{e_i}^x \in \nu(u_i)$ , and  $y \in \cup \nu^{fs}(v_i)$  where  $v_i = \iota(v_{i_1}) \dots \iota(v_{i_n})$ . Thus, there are some  $u \in \nu^{fs}$  and  $v \in \nu^{fs}$  where  $u = \iota(u_1) \dots \iota(u_n)$  and  $x \in \cup \nu^{fs}(u)$ , and  $v \ll \theta \circ f(\iota(v_1) \dots \iota(v_n))$  where  $y \in \cup \nu^{fs}(v)$ . It is obvious that  $\cup \nu^{fs}(u) \cap \cup \nu^{fs}(v) = \tilde{\emptyset}$ . Therefore,  $H'$  is the c.e. set for  $SCpT_2$ .  $\square$

**Example 5.7.** Let  $X = \{x_i, y_i : i \in \mathbb{N}\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  and generated by the following base notation where  $A$  is a non c.e. set,

	$0^i 11$	$0^i 12$	$0^i 21$	$0^i 22$	$0^i 2122$
$i \in A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$\tilde{\emptyset}$
$i \notin A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$y_{E_i}$	$y_{E_i}$	$y_{E_i}$

We extend names to the finite intersections as follows  $\nu(0^i mn) \cap \nu(0^i kl) = \nu(0^i mnkl)$  and the intersection of two basic open sets is empty except for  $\nu(0^i 21) \cap \nu(0^i 22) = \nu(0^i 2122)$ . Thus, the space is computable STS. This space is  $SCpT_2$  as we have a c.e. set  $H_1$  where

$H_1 = \{(\iota(0^i m1)\iota(0^i m2), \iota(0^i n1)\iota(0^i n2)) : i \in \mathbb{N}; m, n \in \{1, 2\}; m \neq n\}$ . Let  $H_2$  be the c.e. set for  $SCuT_2$ , then for

$$i \in A \Rightarrow (0^i 21, 0^i 22) \in H_2,$$

$$i \notin A \Rightarrow (0^i 21, 0^i 22) \notin H_2.$$

Thus,  $A$  must be a c.e. set which is a contradiction. Therefore, the space is not  $SCuT_2$ .

We now give a counterexample for a space that is  $CuT'_1$  but not  $CpT'_1$ .

**Example 5.8.** Let  $X = \{x_i, y_i : i \in \mathbb{N}\}$ ,  $E = \{e_1, e_2\}$  be a parameter set, and  $\tau$  be a STS defined on  $X$  w.r.t.  $E$  and generated by the following base notation where  $A$  is a non c.e. set,

We define names to the finite intersections as follows  $\nu(0^i m) \cap \nu(0^i n) = \nu(0^i mn)$  and the intersection of more than two basic open sets is empty except for  $\nu(0^i 3) \cap \nu(0^i 4) \cap \nu(0^i 6) = \nu(0^i 346)$ .

	$0^i1$	$0^i2$	$0^i3$	$0^i4$	$0^i5$	$0^i6$
$i \in A$	$p_{e_1}^{x_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$\tilde{\emptyset}$	$p_{e_2}^{y_i}$
$i \notin A$	$p_{e_1}^{y_i}$	$p_{e_2}^{y_i}$	$p_{e_1}^{x_i}$	$x_{E_i} \cup p_{e_1}^{y_i}$	$p_{e_2}^{x_i}$	$p_{e_1}^{x_i} \cup p_{e_2}^{y_i}$

Thus, the space is computable STS. This space is  $CuT'_1$  as we have the following c.e. set that satisfies the conditions of  $CuT'_1$ ,

$$H = \{(0^i m, 0^i n) \wedge (0^i 46, 0^i m) : i, j \in \mathbb{N}; m \in \{1, 2, 3, 5\}; n \in \{1, 2, 3, 4\}\}.$$

However, it is not  $CpT'_1$ , as if it was, there would exist a c.e. set  $H'$  that satisfies the conditions of  $CpT'_1$  and hence for,

$$i \in A \Rightarrow (r, s) \in H',$$

$$i \notin A \Rightarrow (r, s) \notin H',$$

where

$$\iota(0^i 1), \iota(0^i 2) \ll r \wedge \iota(0^i 3), \iota(0^i t) \ll s \text{ for } s \in \{4, 6, 46\}.$$

Thus,  $A$  must be a c.e. set which is a contradiction. Therefore, the space is not  $CpT'_1$ .

**Remark 5.9.**  $SCuT_0$  and  $SCpT_0$  are incomparable.

*Proof.* This follows directly from propositions 3.9, 4.7 and examples 5.3 and 5.4.  $\square$

**Remark 5.10.**  $CuT'_i$  and  $CpT'_i$  are incomparable for  $i \in \{0, 1, 2\}$ .

*Proof.* For  $i = 0$ : Use example 5.5 where in which the space is not  $WCuT_0$  and example 5.4, and propositions 3.9, 3.10.

For  $i = 1, 2$ : Use examples 5.5, 5.7, and propositions 3.11, 4.11.  $\square$

**Remark 5.11.**  $WCuT_0$  and  $WCpT_0$  are incomparable.

*Proof.* Use examples 5.5, 4.8 where in the latter example the space is  $WCuT_0$  as we have the following c.e. set,

$$H = \{(0^i mk, 0^i nl) : i, j \in \mathbb{N}; m, n \in \{1, 5\}; k, l \in \{1, 2\}\}.$$

However, it is not  $WCpT_0$  as shown earlier. □

So far we have defined nine computable separation axioms based on soft points and another nine separation axioms based on soft singletons. We also investigated how the ones based on soft points are related and how the other ones that based on soft singletons are related. Counter examples have been provided to show the non-implications between them. Some of them turned out to be equivalent and others turned out to be incomparable. Equivalences between the ones based on soft points exists, however, these equivalences do not exist for their counterparts that based on soft singletons.

In the following figure all relations between computable u-soft and p-soft separation axioms are represented. As seen from the figure, there are some implications between some separation axioms and some other separation axioms turn out to be incomparable.




Figure 3: Relations between computable u-soft and computable p-soft separation axioms.

## References

- [1] ACAR, U., KOYUNCU, F., AND TANAY, B. Soft sets and soft rings. *Computers & Mathematics with Applications* 59, 11 (2010), 3458–3463.
- [2] AKTAŞ, H., AND ÇAĞMAN, N. Soft sets and soft groups. *Information sciences* 177, 13 (2007), 2726–2735.
- [3] AL-SHAMI, T. Comments on some results related to soft separation axioms. *Afrika Matematika* 31, 7 (2020), 1105–1119.
- [4] AL-SHAMI, T. On soft separation axioms and their applications on decision-making problem. *Mathematical Problems in Engineering* 2021 (2021).
- [5] ALI, M. I., FENG, F., LIU, X., MIN, W. K., AND SHABIR, M. On some new operations in soft set theory. *Computers & Mathematics with Applications* 57, 9 (2009), 1547–1553.
- [6] EL-SHAFAEI, M., ABO-ELHAMAYEL, M., AND AL-SHAMI, T. Partial soft separation axioms and soft compact spaces. *Filomat* 32, 13 (2018), 4755–4771.
- [7] EL-SHAFAEI, M., AND AL-SHAMI, T. Applications of partial belong and total non-belong relations on soft separation axioms and decision-making problem. *Computational and Applied Mathematics* 39, 3 (2020), 1–17.
- [8] FENG, F., LI, Y., AND LEOREANU-FOTEA, V. Application of level soft sets in decision making based on interval-valued fuzzy soft sets. *Computers & Mathematics with Applications* 60, 6 (2010), 1756–1767.
- [9] GORZALCZANY, M. B. A method of inference in approximate reasoning based on interval-valued fuzzy sets. *Fuzzy sets and systems* 21, 1 (1987), 1–17.
- [10] JAYNES, E. T. *Probability theory: The logic of science*. Cambridge university press, 2003.
- [11] JUN, Y. B., AND PARK, C. H. Applications of soft sets in ideal theory of bck/bci-algebras. *Information sciences* 178, 11 (2008), 2466–2475.
- [12] L. A., Z. Fuzzy sets. *Information and Control* 8 (1965), 338–353.

- [13] MAJI, P., ROY, A. R., AND BISWAS, R. An application of soft sets in a decision making problem. *Computers & Mathematics with Applications* 44, 8-9 (2002), 1077–1083.
- [14] MOLODTSOV, D. Soft set theory—first results. *Computers & Mathematics with Applications* 37, 4-5 (1999), 19–31.
- [15] NAZMUL, S., AND SAMANTA, S. Neighbourhood properties of soft topological spaces. *Ann. Fuzzy Math. Inform* 6, 1 (2013), 1–15.
- [16] PAWLAK, Z. Rough sets. *International journal of computer & information sciences* 11, 5 (1982), 341–356.
- [17] PEI, D., AND MIAO, D. From soft sets to information systems. In *2005 IEEE international conference on granular computing* (2005), vol. 2, IEEE, pp. 617–621.
- [18] PENG, X., YANG, Y., SONG, J., AND JIANG, Y. Pythagorean fuzzy soft set and its application. *Computer Engineering* 41, 7 (2015), 224–229.
- [19] PEYGHAN, E. About soft topological spaces. *Journal of New Results in Science* 2, 2 (2013), 60–75.
- [20] SHABIR, M., ALI, M. I., AND SHAHEEN, T. Another approach to soft rough sets. *Knowledge-Based Systems* 40 (2013), 72–80.
- [21] SHABIR, M., AND NAZ, M. On soft topological spaces. *Computers & Mathematics with Applications* 61, 7 (2011), 1786–1799.
- [22] TURING, A. M., ET AL. On computable numbers, with an application to the entscheidungsproblem. *J. of Math* 58, 345-363 (1936), 5.
- [23] WEIHRAUCH, K. *Computable analysis: an introduction*. Springer Science & Business Media, 2000.
- [24] WEIHRAUCH, K. Computable separation in topology, from  $t_0$  to  $t_2$ . *J. Univers. Comput. Sci.* 16, 18 (2010), 2733–2753.
- [25] WEIHRAUCH, K. *Computability*, vol. 9. Springer Science & Business Media, 2012.
- [26] WEIHRAUCH, K., AND GRUBBA, T. Elementary computable topology. *J. Univers. Comput. Sci.* 15, 6 (2009), 1381–1422.

- [27] XIAO, Z., GONG, K., AND ZOU, Y. A combined forecasting approach based on fuzzy soft sets. *Journal of Computational and Applied Mathematics* 228, 1 (2009), 326–333.
- [28] YANG, Y., TAN, X., AND MENG, C. The multi-fuzzy soft set and its application in decision making. *Applied Mathematical Modelling* 37, 7 (2013), 4915–4923.