



## Immunisation and Additional Screening Form for LKC**Medicine** students

**PERSONAL PARTICULARS**

Full name (Block letters) : \_\_\_\_\_

NRIC no. / FIN : \_\_\_\_\_ Sex: \_\_\_\_\_ Date of birth: \_\_\_\_\_ Handphone no: \_\_\_\_\_

Please bring along all available immunisation records for the information required below.

1. **Measles, mumps, rubella (MMR) vaccination\***

Dose	Date of MMR vaccination
1	
2	

**OR**, if you received MMR as individual vaccines, or you have serological proof of immunity:

Dose	Date of Measles vaccination
1	
2	

**OR**

Measles antibody level:	
Date:	

Dose	Date of Mumps vaccination
1	
2	

**OR**

Mumps antibody level:	
Date:	

Dose	Date of Rubella vaccination
1	

**OR**

Rubella antibody level:	
Date:	

2. **Chickenpox vaccination#**

Dose	Date of chickenpox vaccination
1	
2	

**OR**

Proof of chickenpox immunity	
Varicella IgG level:	
Date:	

3. **Diphtheria, Pertussis & Tetanus (Tdap) vaccination\***

Dose	Date of last Tdap vaccination
1	

4. **Hepatitis B vaccination & screening test**

Dose	Date of Hepatitis B vaccination	Date of additional Hepatitis B booster / vaccination
1		
2		
3		

HBs Ag: \_\_\_\_\_ Anti-HBs antibody: \_\_\_\_\_ Date: \_\_\_\_\_

Students with HBs Ab < 10 mIU/ml will need further Hepatitis B immunisation followed by repeat anti-HBs antibody level testing.

Post-booster immunisation Anti-HBs antibody: \_\_\_\_\_ Date: \_\_\_\_\_

5. **Hepatitis C test^**

Anti-HCV IgG test: \_\_\_\_\_ Date: \_\_\_\_\_

6. **HIV test**

Anti-HIV antibody: \_\_\_\_\_ Date: \_\_\_\_\_

\_\_\_\_\_  
Physician's Signature, Name & Professional Qualification

\_\_\_\_\_  
Clinic stamp

Notes:

+ All medical students must receive 2 doses of Measles, Mumps & Rubella (MMR) vaccine, or separate measles, mumps and rubella vaccine or, are able to demonstrate serological proof of immunity.

# Chickenpox vaccination is given in two doses, at least four weeks apart.

\* Last dose of Diphtheria, Pertussis & Tetanus (Tdap) vaccine is usually given at age 11-12 in Singapore. Documented proof of vaccination with Tdap or Td within the last 10 years is acceptable as evidence of immunity.

If vaccination was more than 10 years ago, a booster dose of Tdap is required.

^ Positive serology (anti-HCV IgG positive) must be followed up by HCV RNA test. Individuals who test anti-HCV IgG positive but HCV RNA test negative can be considered to be non-carriers. They need to obtain a medical report from their certifying physician that they have cleared their Hepatitis C infection.